

**The following resources related to this article are available online at  
[www.sciencemag.org](http://www.sciencemag.org) (this information is current as of December 29, 2009):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/325/5947/1541>

**Supporting Online Material** can be found at:

<http://www.sciencemag.org/cgi/content/full/325/5947/1541/DC1>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/cgi/content/full/325/5947/1541#related-content>

This article **cites 25 articles**, 8 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/325/5947/1541#otherarticles>

This article has been **cited by** 1 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/cgi/content/full/325/5947/1541#otherarticles>

This article appears in the following **subject collections**:

Microbiology

<http://www.sciencemag.org/cgi/collection/microbio>

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

less of the level of taxonomic resolution. In this regard, genetic analyses have shown the existence of widespread, although disjunct, distribution patterns for several cryptic species of the marine diatom *Skeletonema* (23). Similar results have been reported for other microbial plankton groups, such as picoeukaryote algae and foraminifera (17, 24). Alternatively, our results could reflect the convergence or parallelism of morphological traits among genetically unrelated taxa in disjunct oceanic regions.

Despite enormous environmental variability linked to glacial-interglacial climates of the Pleistocene, our analysis reveals that marine diatom communities have evolved slowly through gradual changes over the past 1.5 My of Earth's history (Fig. 2 and table S2). These patterns of community stability for extensive periods of geological time are probably associated with the great dispersal ability of marine diatoms (25, 26) and highlight the potential of microbial plankton communities for recovering from past and future climatic variations. This conclusion implies that there are few or no biogeographical traces of historical climate change in contemporary communities of marine diatoms.

Models of evolution of species commonly assume that tectonic barriers and water mass fronts act as effective isolating mechanisms (9). This is a necessary condition that precedes the delineation of biogeographic provinces *sensu stricto* (6, 9) and controls the development of global species richness. Our analysis, however, indicates that, even at the largest spatial scale, the geographic distribution of marine planktonic diatoms does

not seem to be limited by dispersal. These results, together with recent genetic evidence for high rates of inter- and intra-oceanic gene flow in planktonic protists and widespread oceanic distributions of cryptic "sibling" species (17, 23, 27), suggest that the geographic isolation of marine diatoms cannot be maintained for long periods. Our results strongly support the hypothesis that environmental selection rather than dispersal dominates diatom community structure. To the extent that marine diatoms are a model microbial taxonomic group, our results imply that the biodiversity and macroevolutionary patterns at the microbial level fundamentally differ from those of macroscopic animals and plants, negating the idea that all living things follow similar ecological and evolutionary rules (6).

#### References and Notes

1. L. G. M. Baas-Becking, *Geobiologie of Inleiding Tot de Milieukunde* (Van Stockum & Zoon, The Hague, Netherlands, 1934).
2. B. J. Finlay, *Science* **296**, 1061 (2002).
3. T. Fenchel, B. J. Finlay, *Philos. Trans. R. Soc. London Ser. B Biol. Sci.* **361**, 1965 (2006).
4. R. J. Whitaker, D. W. Grogan, J. W. Taylor, *Science* **301**, 976 (2003).
5. B. P. Hedlund, J. T. Staley, in *Microbial Diversity and Bioprospecting*, A. T. Bull, Ed. (ASM Press, Washington DC, 2004), pp. 225–231.
6. J. B. Hughes-Martiny *et al.*, *Nat. Rev. Microbiol.* **4**, 102 (2006).
7. R. J. Telford, V. Vandvik, H. J. B. Birks, *Science* **312**, 1015 (2006).
8. S. P. Hubbell, *The Unified Neutral Theory of Biodiversity and Biogeography* (Princeton Univ. Press, Princeton, NJ, 2001).
9. M. V. Lomolino, B. R. Riddle, J. H. Brown, *Biogeography* (Sinauer, Sunderland, MA, 2006).
10. W. J. Lee, D. J. Patterson, *Protist* **149**, 229 (1998).

11. J. J. Cerverto, *Palaeontol. Electronica* **2**, 1 (1999).
12. Materials and methods are available as supporting online material on Science Online.
13. A. R. Longhurst, S. Sathiyendranath, T. Platt, C. Caverhill, *J. Plankton Res.* **17**, 1245 (1995).
14. R. F. Strzepak, P. J. Harrison, *Nature* **431**, 689 (2004).
15. J. Soinenen, R. Paavola, T. Muotka, *Ecography* **27**, 330 (2004).
16. W. Vyverman *et al.*, *Ecology* **88**, 1924 (2007).
17. C. de Vargas, R. Norris, L. Zaninetti, S. W. Gidd, J. Pawlowski, *Proc. Natl. Acad. Sci. U.S.A.* **96**, 2864 (1999).
18. R. D. Norris, in *Reconstructing Ocean History: A Window into the Future*. F. Abrantes, A. Mix, Eds. (Kluwer/Plenum, New York, 1999), pp. 173–193.
19. P. F. Sexton, R. D. Norris, *Geology* **36**, 899 (2008).
20. D. J. McGillicuddy *et al.*, *Science* **316**, 1021 (2007).
21. D. G. Mann, *Phycologia* **38**, 437 (1999).
22. P. Vanormelingen, E. Verleyen, W. Vyverman, *Biodivers. Conserv.* **17**, 393 (2008).
23. W. H. Kooistra *et al.*, *Protist* **159**, 177 (2008).
24. J. Slapeta, P. López-García, D. Moreira, *Mol. Biol. Evol.* **23**, 23 (2005).
25. W. A. DiMichele *et al.*, *Annu. Rev. Ecol. Syst.* **35**, 285 (2004).
26. C. de Vargas, A. G. Saez, L. Medlin, H. R. Thierstein, in *Coccolithophores: From Molecular Processes to Global Impact*, H. R. Thierstein, J. R. Young, Eds. (Springer-Verlag, Berlin Heidelberg, 2004), pp. 251–298.
27. K. F. Darling *et al.*, *Nature* **405**, 43 (2000).
28. We thank all the people who contributed to the collection and management of data and C. de Vargas, J. Grasse, M. Katz, and P. Morin for comments on the manuscript. The Neptune database is hosted by Chronos ([www.chronos.org](http://www.chronos.org)). This work was supported by a Marie Curie International Fellowship within the 6th European Community framework Program (P.C.).

#### Supporting Online Material

[www.sciencemag.org/cgi/content/full/325/5947/1539/DC1](http://www.sciencemag.org/cgi/content/full/325/5947/1539/DC1)  
 Materials and Methods

Fig. S1  
 Tables S1 and S2  
 References

27 March 2009; accepted 30 July 2009  
 10.1126/science.1174159

## A Constant Flux of Diverse Thermophilic Bacteria into the Cold Arctic Seabed

Casey Hubert,<sup>1\*</sup> Alexander Loy,<sup>2</sup> Maren Nickel,<sup>1</sup> Carol Arnosti,<sup>3</sup> Christian Baranyi,<sup>2</sup> Volker Brüchert,<sup>1†</sup> Timothy Ferdelman,<sup>1</sup> Kai Finster,<sup>4</sup> Flemming Mønsted Christensen,<sup>4,5</sup> Júlia Rosa de Rezende,<sup>1,5</sup> Verona Vandieken,<sup>1‡</sup> Bo Barker Jørgensen<sup>1,5</sup>

Microorganisms have been repeatedly discovered in environments that do not support their metabolic activity. Identifying and quantifying these misplaced organisms can reveal dispersal mechanisms that shape natural microbial diversity. Using endospore germination experiments, we estimated a stable supply of thermophilic bacteria into permanently cold Arctic marine sediment at a rate exceeding 10<sup>8</sup> spores per square meter per year. These metabolically and phylogenetically diverse *Firmicutes* show no detectable activity at cold in situ temperatures but rapidly mineralize organic matter by hydrolysis, fermentation, and sulfate reduction upon induction at 50°C. The closest relatives to these bacteria come from warm subsurface petroleum reservoir and ocean crust ecosystems, suggesting that seabed fluid flow from these environments is delivering thermophiles to the cold ocean. **These transport pathways may broadly influence microbial community composition in the marine environment.**

Microbial diversity surveys have revealed that species richness is determined by many low-abundance taxa—the so-called rare biosphere (1–3). In the ocean, certain

members of this relatively unexplored biosphere comprise a dormant microbial “seed bank” that can be transported passively over great distances (1). Quantitatively tracking the migration of in-

dicator taxa can highlight key factors that influence patterns of biogeography and may help evaluate the extent to which microorganisms exhibit a cosmopolitan distribution (4). Endospore germination allows certain bacteria to persist as dormant cells in hostile environments, explaining discoveries of viable thermophilic *Firmicutes* in inhospitably cold habitats (5–10). Quantitative studies of this phenomenon are scarce, and the origin and distribution of thermophiles in cold environments remain enigmatic (6–11). Thermophilic sporulating taxa such as certain *Desulfotomaculum* spp. may constitute only 0.001% of marine microbial populations (8, 12). Like the rare taxa, spores are less prone to viral lysis or predation, and are not detected by traditional diversity surveys (1, 2). A spore-forming *Desulfotomaculum* strain that can only grow between 26°C and 47°C was recently isolated from permanently cold Svalbard fjord sediment in the Arctic (10). The present study assessed thermophile diversity, abundance, and distribution in Svalbard sediments to reveal insights into mechanisms governing biogeography in the marine environment.

Pristine sediment was sampled from **Smeerenburgfjorden (80°N; fig. S1)** and incubated over an experimental temperature range (13), which revealed two distinct sulfate-reduction regimes


(Fig. 1A). The first has a temperature optimum ( $T_{opt}$ ) of 22°C and a maximum ( $T_{max}$ ) of 32°C, consistent with earlier studies in Svalbard sediments (14, 15) where the in situ temperature is -2° to +4°C year-round. This temperature-activity profile is characteristic of psychrophilic sulfate-reducing bacteria (SRB) that are well adapted to cold conditions (14, 16), which explains their activity and abundance (up to 16%) in this environment (17). A second  $T_{opt}$  of 56°C (Fig. 1) indicates a previously unrecognized thermophilic community with a  $T_{max}$  above 60°C. Pasteurization at 80°C killed the psychrophiles but did not adversely affect the thermophiles (Fig. 1B), indicating that the latter exist as endospores in situ and only germinate after heating. This was supported by successful polymerase chain reaction targeting the spore-forming SRB genus *Desulfotomaculum* only if the sediment was incubated at 50°C before DNA extraction.

Psychrophilic and thermophilic communities were investigated further by incubating homogenized surface sediment at 50°C (13). Sulfate reduction rates (SRR) quickly dropped to below the detection limit (Fig. 2A) due to stimulation and subsequent death of vegetative psychrophiles as the sediment warmed up (compare to Fig. 1A). Thermophilic SRR increased exponentially between 20 and 96 hours. The transition from SRR below detection (up to 16 hours) to the onset of the exponential phase (at 20 hours) suggests a lag phase during which conditions became favorable for germination of thermophilic SRB spores. Endospore germination requires appropriate nutrients and substrates (18) such as volatile fatty acids (VFA) that are produced by microbial fermentation. VFA are typical electron donors for SRB and were generated rapidly during incubation at 50°C (Fig. 2B), creating conditions suitable for thermophilic sulfate reduction.


Several lines of evidence demonstrate that the VFA production was biologically mediated and not due to thermal alteration of the sediment organic matter. VFA did not accumulate in autoclaved sediment incubated in parallel at 50°C (concentrations never exceeded 0.3 mM). Amendment with the fluorescently labeled polysaccharides pullulan or arabinogalactan revealed extracellular enzymatic hydrolysis in sediment

incubated at 50°C, whereas no hydrolysis was detected in sediment-free controls. Clone library analyses of 16S ribosomal RNA (rRNA) genes revealed enrichment at 50°C of different bacterial groups related to *Desulfotomaculum*, *Caminicella*, and *Caloranaerobacter-Clostridiisalibacter-Thermohalobacter* lineages within the *Firmicutes* phylum (Fig. 3) [see supporting online material (SOM) and table S3]. The latter lineages are

**Fig. 1.** Temperature-gradient incubations. SRR were measured in Smeerenburgfjorden sediment samples incubated between 0° and 76°C. Distinct SRB populations were active between 0° and 32°C, and between 41° and 62°C. Replicates were either untreated (A) or pasteurized for 1 hour at 80°C (B) before incubation. Open and closed symbols correspond to experiments below and above 35°C, respectively. Incubations in the thermophilic range resulted in much higher SRR (fig. S3); therefore, data are plotted as the percentage of the maximum SRR in (A) at either  $T_{opt}$ . Pasteurization killed the psychrophilic community and resulted in slightly higher activity by the thermophilic community [115% at the  $T_{opt}$  (B)].



**Fig. 2.** Sediment incubation at 50°C. SRR (A) and concentrations of VFA (B) in Smeerenburgfjorden sediment (0- to 3-cm depth) incubated at 50°C. SRR represent 4- to 8-hour incubations with  $^{35}\text{SO}_4^{2-}$  radiotracer. Open symbols correspond to the psychrophilic SRB community that was killed as the temperature increased to beyond their  $T_{max}$  (compare to Fig. 1A). SRR were below detection during the 10 to 16 hours time interval ( $n = 2$ ; vertical bar: standard error). Sulfate reduction by 20 to 24 hours indicates germination of *Desulfotomaculum* spores, which subsequently catalyzed an exponential increase in SRR. These data constrain the earliest onset of the exponential phase to 16 hours, as indicated by the dashed extension of the exponential trendline. The exponential phase corresponded to the consumption of butyrate and propionate, which increased steeply, together with acetate (B), before the onset of thermophilic sulfate reduction.



<sup>1</sup>Biogeochemistry Group, Max Planck Institute for Marine Microbiology, Celsiusstrasse 1, D-28359 Bremen, Germany.

<sup>2</sup>Department of Microbial Ecology, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria. <sup>3</sup>Department of Marine Sciences, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-3300, USA. <sup>4</sup>Department of Biological Sciences—Microbiology Section, Aarhus University, Ny Munkegade, Building 1540, DK-8000 Aarhus C, Denmark. <sup>5</sup>Center for Geomicrobiology, Department of Biological Sciences, Aarhus University, Ny Munkegade, Building 1535, DK-8000 Aarhus C, Denmark.

\*To whom correspondence should be addressed. E-mail: chubert@mpi-bremen.de

†Present address: Department of Geology and Geochemistry, Stockholm University, Svante Arrhenius Väg 8C, 106 91 Stockholm, Sweden.

‡Present address: Institute of Biology, University of Southern Denmark, Campusvej 55, 5320 Odense M Denmark.

represented by thermophilic anaerobes that convert carbohydrates and proteins to VFA, such as *Caminicella sporogenes* (19). Hydrolysis and fermentation at 50°C were therefore likely due to thermophilic spores that germinated at high temperature in the presence of complex natural organic substrates in the sediment.

VFA production at 50°C stimulated growth of a thermophilic *Desulfotomaculum* population

(Fig. 3), resulting in an exponential increase in SRR (Fig. 2A). The *Desulfotomaculum* cell density at the onset of sulfate reduction can be estimated by dividing the bulk rate ( $1.3 \text{ nmol cm}^{-3} \text{ hour}^{-1}$  at 16 hours based on the exponential function in Fig. 2A) by a mean cell-specific sulfate reduction rate of  $2.0 \text{ fmol cell}^{-1} \text{ hour}^{-1}$  [based on 33 pure cultures during exponential growth (20), and representative of thermophilic *Desulfotomaculum* spp.; see SOM]. This calculation estimates a population of  $6.3 \times 10^5 \text{ SRB cm}^{-3}$  at 16 hours, which corresponds to the in situ spore density given that SRB growth did not occur before 16 hours (Fig. 2A).

Exponential increases in SRR were also measured across a series of intact sediment cores (0- to 23-cm depth) incubated at  $50^\circ\text{C}$  (13). These data indicate similar numbers of thermophilic *Desulfotomaculum* spores, on the order of  $10^5 \text{ cm}^{-3}$ , at all depths (fig. S2). Thermophiles thus constitute a stable component of the rare biosphere ( $\sim 0.01\%$ ) in this Arctic marine habitat (17). Abundant taxa in Smeerenburgfjorden sediment are psychrophilic *Cytophaga*, *Flavobacteria*, *Gammaproteobacteria*, and *Deltaproteobacteria*


(17) that catalyze hydrolysis, fermentation, and sulfate reduction during degradation of organic matter (15, 21). We induced the same processes at  $50^\circ\text{C}$ , which stimulated SRR about 30-fold higher than in sediment incubated at  $4^\circ\text{C}$  (fig. S3). It is interesting that the thermophilic bacterial community has metabolic potential that mirrors the dominant metabolic processes occurring in situ. However, these thermophilic spores make no contribution to local biogeochemical cycling (Fig. 1). Our experience with thermophiles cultivated from Smeerenburgfjorden that have no detectable metabolic activity below  $25^\circ\text{C}$  supports this. Thermophilic spores must therefore be supplied externally and can be considered akin to particles accumulating in the sediment.

To estimate the particle flux of thermophilic spores into the sediment, we determined an age model by measuring the  $^{210}\text{Pb}$  activity as a function of depth (see SOM). This revealed a sediment accumulation rate of  $0.19 \text{ cm year}^{-1}$  in Smeerenburgfjorden (figs. S2 and S4). Given a stable abundance of about  $10^5$  spores  $\text{cm}^{-3}$  with depth (fig. S2), the constant supply of thermophiles to this site is  $2 \times 10^8 \text{ m}^{-2} \text{ year}^{-1}$ . Other

findings along the west coast of Svalbard have comparable thermophilic SRB densities in surface sediments (see SOM; fig. S1). The fjords and the adjacent coastal shelf in this area cover about  $1000 \text{ km}^2$  (fig. S1), hence an extrapolation of our estimate corresponds to an annual supply of  $\sim 10^{17}$  thermophiles in this Arctic region. These large numbers highlight not merely the occurrence of thermophiles in the Arctic but rather their unexpected quantity and the consistency of their flux. The warm, anaerobic environment where these bacteria originate must have sufficient geographic distribution, magnitude, and source strength to support the population sizes indicated by our data.

A limited number of marine habitats meet these criteria. Deeply buried sediments hundreds of meters below the sea floor represent warm habitats for anaerobic microbes (22). Advective flow of hydrocarbons or other fluids from these sediments can penetrate the sea floor (e.g., at mud volcanoes) and transport microscopic particles or cells up into the cold ocean (23). Sea-floor pockmarks and active cold seeps are known around west Svalbard (24) (fig. S1) and are often

**Fig. 3.** Phylogenetic analysis of putative spore-forming thermophilic bacteria in Svalbard sediment. Clone libraries constructed before and after incubation at  $50^\circ\text{C}$  differed significantly due to the enrichment of thermophilic *Firmicutes* (see SOM, tables S2 and S3). Dominant phylotypes (i.e.,  $>5\%$  of clones) from the  $50^\circ\text{C}$  sediment library are indicated in bold-face with relative abundances shown in parentheses. These phylotypes were not detected before  $50^\circ\text{C}$  incubation. The consensus tree of 16S rRNA gene sequences of selected *Firmicutes* includes next relatives for each phylotype (top 10) plus additional type strains (for *Desulfotomaculum*). Habitats of origin for closest relatives are indicated. Filled and open circles indicate lineages with  $>90\%$  and 80 to  $90\%$  parsimony bootstrap support (100 resamplings), respectively. The scale bar indicates 5% sequence divergence as estimated from maximum-likelihood analysis.



associated with deep gas or oil-bearing deposits (23). **16S rRNA gene sequence comparisons revealed that taxa enriched in our 50°C experiments (Fig. 3) are most closely related to bacteria from subsurface petroleum reservoirs or oil production facilities (94 to 96% similarity; table S3).**

Another source of thermophiles could be nearby mid-ocean ridge spreading centers (fig. S1). Large volumes of fluids circulating through ocean crust (25) could transport cells away from warm **anoxic niches** in this seafloor habitat (26) and suspend them in abyssal currents. **The closest relatives to the Arctic thermophiles also include an anaerobic thermophile isolated from deep, hot crustal fluid (94% similarity; table S3) (27).**

Petroleum-bearing sediments and fractured ocean crust both host anaerobic heterotrophic microbial communities (26, 28). Areas of discharge connecting these habitats to the water column are widespread, and both processes expel large volumes of fluid into the oceans (23, 25). A combination of different point sources could explain the diversity and distribution of thermophilic taxa in Arctic sediments (Fig. 3 and fig. S1). Although our observations suggest that seabed fluid flow governs the biogeography of thermophilic spore formers, these passive dispersal mechanisms are unlikely to act only on these particular bacteria. Permeable conduits through sediments and ocean crust pass through several microbial niches with changing local temperature

and geochemistry (22, 25, 26). Widespread seeding of the oceans by geofluids from deep biosphere habitats may therefore contribute broadly to the high microbial diversity observed in the marine environment.

#### References and Notes

- C. Pedr s-Ali , *Trends Microbiol.* **14**, 257 (2006).
- M. L. Sogin *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 12115 (2006).
- J. A. Huber *et al.*, *Science* **318**, 97 (2007).
- L. G. M. Baas-Becking, *Geobiologie of inleiding Tot de Milieukunde* (Van Stockum & Zoon, The Hague, Netherlands, 1934).
- A. A. Egorova, *C. R. (Dokl.) Acad. Sci. USSR* **19**, 649 (1938).
- R. H. McBee, V. H. McBee, *J. Bacteriol.* **71**, 182 (1956).
- J. W. Bartholomew, G. Paik, *J. Bacteriol.* **92**, 635 (1966).
- M. F. Isaksen, F. Bak, B. B. J rgensen, *FEMS Microbiol. Ecol.* **14**, 1 (1994).
- Y. J. Lee *et al.*, *Extremophiles* **9**, 375 (2005).
- V. Vandieken, C. Knoblauch, B. B. J rgensen, *Int. J. Syst. Evol. Microbiol.* **56**, 687 (2006).
- K. O. Stetter *et al.*, *Nature* **365**, 743 (1993).
- K. Sahn, B. J. MacGregor, B. B. J rgensen, D. A. Stahl, *Environ. Microbiol.* **1**, 65 (1999).
- Materials and methods are available as supporting material on Science Online.
- J. Sagemann, B. B. J rgensen, O. Greif, *Geomicrobiol. J.* **15**, 85 (1998).
- N. Finke, B. B. J rgensen, *ISME J.* **2**, 815 (2008).
- C. Knoblauch, B. B. J rgensen, *Environ. Microbiol.* **1**, 457 (1999).
- K. Ravensschlag, K. Sahn, R. Amann, *Appl. Environ. Microbiol.* **67**, 387 (2001).
- P. Setlow, *Curr. Opin. Microbiol.* **6**, 550 (2003).
- K. Alain *et al.*, *Int. J. Syst. Evol. Microbiol.* **52**, 1621 (2002).

本页已使用福昕阅读器进行编辑。  
福昕软件(C)2005-2007, 版权所有, 仅供试用。

- J. A. Huber, V. Br uchert, K. S. Habicht, J. Kuever, *Appl. Environ. Microbiol.* **67**, 888 (2001).
- C. Arnosti, B. B. J rgensen, *Geomicrobiol. J.* **23**, 551 (2006).
- R. J. Parkes *et al.*, *Org. Geochem.* **38**, 845 (2007).
- A. G. Judd, M. Hovland, *Seabed Fluid Flow: The Impact of Geology, Biology and the Marine Environment* (Cambridge Univ. Press, Cambridge, 2007).
- E. Damm, A. Mackensen, G. Bud us, E. Faber, C. Hanfland, *Cont. Shelf Res.* **25**, 1453 (2005).
- M. Hutnak *et al.*, *Nat. Geosci.* **1**, 611 (2008).
- J. P. Cowen, *Res. Microbiol.* **155**, 497 (2004).
- J. A. Huber, P. Johnson, D. A. Butterfield, J. A. Baross, *Environ. Microbiol.* **8**, 88 (2006).
- I. M. Head, D. M. Jones, S. R. Larter, *Nature* **426**, 324 (2003).
- We thank S. Henningsen, J. Mortensen, K. Barker, and A. Steen for assistance aboard R/V *Farm* and the Alfred Wegener Institute for kindly providing laboratory space in Ny  lesund, Svalbard (project KOP56; RIS ID: 3298). We are grateful to K. Imhoff, M. Meyer, and A. Schipper for technical assistance and to A. Judd, M. Holtappels, N. Finke, and W. Bach for valuable discussions. This work was supported by the Natural Sciences and Engineering Research Council of Canada (C.H.), the Austrian Science Fund (P20185-B17; A.L.), the U.S. National Science Foundation (OCE-0323975 and OCE-0848703; C.A.), and the Max Planck Society. Nucleotide sequences have been deposited in the European Molecular Biology Laboratory database under accession numbers FN396615 to FN396795.

#### Supporting Online Material

www.sciencemag.org/cgi/content/full/325/5947/1541/DC1

Materials and Methods

SOM Text

Figs. S1 to S4

Tables S1 to S3

24 March 2009; accepted 20 July 2009

10.1126/science.1174012

## Three-Dimensional Structural View of the Central Metabolic Network of *Thermotoga maritima*

Ying Zhang,<sup>1\*</sup> Ines Thiele,<sup>2,\*†</sup> Dana Weekes,<sup>3</sup> Zhanwen Li,<sup>1</sup> Lukasz Jaroszewski,<sup>3</sup> Krzysztof Ginalski,<sup>4</sup> Ashley M. Deacon,<sup>5</sup> John Wooley,<sup>6</sup> Scott A. Lesley,<sup>7</sup> Ian A. Wilson,<sup>8</sup> Bernhard Palsson,<sup>2</sup> Andrei Osterman,<sup>9</sup> Adam Godzik<sup>1,3,6‡</sup>

Metabolic pathways have traditionally been described in terms of biochemical reactions and metabolites. With the use of structural genomics and systems biology, we generated a three-dimensional reconstruction of the central metabolic network of the bacterium *Thermotoga maritima*. The network encompassed 478 proteins, of which 120 were determined by experiment and 358 were modeled. Structural analysis revealed that proteins forming the network are dominated by a small number (only 182) of basic shapes (folds) performing diverse but mostly related functions. Most of these folds are already present in the essential core (~30%) of the network, and its expansion by nonessential proteins is achieved with relatively few additional folds. Thus, integration of structural data with networks analysis generates insight into the function, mechanism, and evolution of biological networks.

The advent of genome sequencing has enabled development of computational and experimental tools to investigate complete biological systems, but it has also highlighted the difficulty in integrating complex information for the hundreds to thousands of different molecules that compose even the smallest biological networks. Such integration presents many challenges, especially when assembling data from

diverse fields, such as biochemistry and structural biology, that use different operational languages and conceptual frameworks. Biochemistry has traditionally focused on individual reactions and pathways, but recent advances in genomics have led to more rapid growth in the reconstruction and modeling of metabolic networks on a genome-wide scale (1–3). Thus, biochemical reactions, pathways, and networks can now be described in

the context of entire cells, thereby enabling more realistic simulations of the behavior of metabolic networks in a growing number of organisms (4–7). Nevertheless, metabolism is still generally defined in terms of the chemical names and identity of substrates, products, and reactions. It does not explicitly consider the three-dimensional structures of its components, although such knowledge is required for a comprehensive understanding, not only of the individual reactions, but more importantly, of metabolic networks as a whole. Without such knowledge, we cannot rigorously define enzyme mechanisms or

<sup>1</sup>Joint Center for Molecular Modeling (JCMM), Burnham Institute for Medical Research, La Jolla, CA 92037, USA. <sup>2</sup>Department of Bioengineering, University of California at San Diego, La Jolla, CA 92093–0412, USA. <sup>3</sup>Joint Center for Structural Genomics (JCSG), Bioinformatics Core, Burnham Institute for Medical Research, La Jolla, CA 92037, USA. <sup>4</sup>Interdisciplinary Centre for Mathematical and Computational Modelling, Warsaw University, Warsaw, Poland. <sup>5</sup>JCSG, Structure Determination Core, Stanford Synchrotron Radiation Lightsource, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA. <sup>6</sup>JCSG, Bioinformatics Core, University of California at San Diego, La Jolla, CA 92093, USA. <sup>7</sup>JCSG, Crystallography Core, Genomics Institute of the Novartis Research Foundation, San Diego, CA 92121, USA. <sup>8</sup>JCSG, The Scripps Research Institute, La Jolla, CA 92037, USA. <sup>9</sup>Burnham Institute for Medical Research, La Jolla, CA 92037, USA.

\*These authors contributed equally to this work.

†Present address: Center of Systems Biology, University of Iceland, IS-101 Reykjavik, Iceland.

‡To whom correspondence should be addressed. E-mail: adam@burnham.org