

# First insights into the biodiversity and biogeography of the Southern Ocean deep sea


Angelika Brandt<sup>1</sup>, Andrew J. Gooday<sup>2</sup>, Simone N. Brandão<sup>1</sup>, Saskia Brix<sup>1</sup>, Wiebke Brökeland<sup>1</sup>, Tomas Cedhagen<sup>3</sup>, Madhumita Choudhury<sup>1</sup>, Nils Cornelius<sup>2</sup>, Bruno Danis<sup>4</sup>, Ilse De Mesel<sup>5</sup>, Robert J. Diaz<sup>6</sup>, David C. Gillan<sup>7</sup>, Brigitte Ebbe<sup>8</sup>, John A. Howe<sup>9</sup>, Dorte Janussen<sup>10</sup>, Stefanie Kaiser<sup>1</sup>, Katrin Linse<sup>11</sup>, Marina Malyutina<sup>12</sup>, Jan Pawlowski<sup>13</sup>, Michael Raupach<sup>14</sup> & Ann Vanreusel<sup>5</sup>

Shallow marine benthic communities around Antarctica show high levels of endemism, gigantism, slow growth, longevity and late maturity, as well as adaptive radiations that have generated considerable biodiversity in some taxa<sup>1</sup>. The deeper parts of the Southern Ocean exhibit some unique environmental features, including a very deep continental shelf<sup>2</sup> and a weakly stratified water column, and are the source for much of the deep water in the world ocean. These features suggest that deep-sea faunas around the Antarctic may be related both to adjacent shelf communities and to those in other oceans. Unlike shallow-water Antarctic benthic communities, however, little is known about life in this vast deep-sea region<sup>2,3</sup>. Here, we report new data from recent sampling expeditions in the deep Weddell Sea and adjacent areas (748–6,348 m water depth) that reveal high levels of new biodiversity; for example, 674 isopods species, of which 585 were new to science. Bathymetric and biogeographic trends varied between taxa. In groups such as the isopods and polychaetes, slope assemblages included species that have invaded from the shelf. In other taxa, the shelf and slope assemblages were more distinct. Abyssal faunas tended to have stronger links to other oceans, particularly the Atlantic, but mainly in taxa with good dispersal capabilities, such as the Foraminifera. The isopods, ostracods and nematodes, which are poor dispersers, include many species currently known only from the Southern Ocean. Our findings challenge suggestions that deep-sea diversity is depressed in the Southern Ocean and provide a basis for exploring the evolutionary significance of the varied biogeographic patterns observed in this remote environment.

Although animal communities inhabiting shallow marine benthic environments around Antarctica, notably parts of McMurdo Sound and the Peninsula region, are well known, there have been few studies of the deep-water faunas in the adjacent Southern Ocean. The ANDEEP (Antarctic benthic deep-sea biodiversity: colonization history and recent community patterns) project was designed to fill this knowledge vacuum<sup>4</sup>. Between 2002 and 2005, we undertook three expeditions in the deep Weddell Sea and adjacent areas aboard the German research vessel *Polarstern*. Biological collections and data on environmental and seafloor characteristics were obtained from diverse settings, including continental slope, rise, abyssal plain,

trench floor, channel levees and adjacent to fracture zones, and from water depths between 774 and 6,348 m (Fig. 1; Supplementary Discussion; Supplementary Tables 1, 2; Supplementary Figs 1, 2). This material has greatly improved our knowledge of the biodiversity of benthic communities in the deep Southern Ocean and enabled us to test ideas about large-scale biogeographic and other macroecological patterns<sup>5</sup> among deep-water faunas.

The richness of species inhabiting deep-sea sediments was first demonstrated during the 1960s<sup>6,7</sup>. Since then, scientists have sought to understand the mechanisms that maintain these high levels of benthic diversity at local<sup>8</sup> and regional<sup>9</sup> scales. We analysed species assemblages in ANDEEP samples across a range of taxonomic groups, representing the meiofauna, macrofauna and megafauna (Fig. 2) and found substantial levels of unrecorded biodiversity (see Supplementary Material). The Foraminifera were represented by 158 live species, including 72 monothalamous species, most of them undescribed. The nematodes belonged to typical cosmopolitan deep-sea genera (Supplementary Fig. 3), but more than half of the 57




**Figure 1 | Station map.** Stations from ANDEEP I (2002) red circles, ANDEEP II (2002) yellow circles and ANDEEP III (2005) green circles. The blue line indicates the Polar Front.

<sup>1</sup>Zoological Museum Hamburg, Martin-Luther-King-Platz 3, 20146 Hamburg, Germany. <sup>2</sup>National Oceanography Centre, Southampton, European Way, Southampton SO14 3ZH, UK. <sup>3</sup>Department of Marine Ecology, University of Århus, Finlandsgade 14, 8200 Århus N, Denmark. <sup>4</sup>Royal Belgian Institute of Natural Sciences, 29 Rue Vautier, 1000 Brussels, Belgium. <sup>5</sup>Biology Department, Marine Biology Section, Ghent University, Krijgslaan 281/S8, 9000 Ghent, Belgium. <sup>6</sup>Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, Virginia 23062, USA. <sup>7</sup>Marine Biology Laboratory, CP160/15, Université Libre de Bruxelles, 50 Avenue Roosevelt, 1050 Brussels, Belgium. <sup>8</sup>Forschungsinstitut Senckenberg, DZMB-CeDAMar, c/o Forschungsmuseum König, Adenauerallee 160, 53113 Bonn, Germany. <sup>9</sup>Scottish Association for Marine Science, Dunstaffnage Marine Laboratory, Dunbeg, Oban, Argyll PA37 1QA, UK. <sup>10</sup>Forschung und Naturmuseum Senckenberg, Senckenberganlage 25, 60325 Frankfurt am Main, Germany. <sup>11</sup>British Antarctic Survey, Natural Environmental Research Council, High Cross, Madingley Road, Cambridge CB3 0ET, UK. <sup>12</sup>Institute of Marine Biology, FEB RAS, Palchevskogo, 17, 690041 Vladivostok, Russia. <sup>13</sup>Department of Zoology & Animal Biology, University of Geneva, 30 Quai Ernest Ansermet, CH 1211 Genève 4, Switzerland. <sup>14</sup>Ruhr-Universität Bochum, Universitätsstrasse 150, 44780 Bochum, Germany.

species recognized in selected genera were new to science. At least 100 ostracod species were distinguished, and >70% of them were new. Among the macrofauna, the isopods were astonishingly diverse with 674 species identified among >13,000 specimens examined, compared with 371 species reported from the entire Antarctic continental shelf<sup>10</sup>. Most (86%) of the isopod species were undescribed and are presently known only from the Southern Ocean. More than 200 polychaete species were recognized, 81 of them previously unknown. Our samples yielded 160 species of shelled gastropods and bivalves compared with 279 species on the shelf (<1,000 m)<sup>11</sup>. Around 40% of species were confined to abyssal depths. Among the megafauna, 76 species of sponges were recognized, 17 of them were undescribed and 37 new for the Southern Ocean (Supplementary Table 3).


Whether shelf faunas have colonized the deep ocean, or vice versa, is a central issue for understanding faunal patterns in the Southern Ocean. Prior to ANDEEP, most of the information on bathymetric distributions around Antarctica concerned shelf and upper-slope faunas<sup>10</sup>. A comparison of species distributions on continental shelves and slopes around Europe and Antarctica revealed significantly wider depth ranges off Antarctica for bivalves, gastropods,



**Figure 2 | Selected important ANDEEP taxa.** **a**, A new species of komokiacean (*Ipoa* sp. nov.) from the Weddell Sea abyssal plain. **b**, The carnivorous sponge *Chondrocladia* from the Weddell Sea abyssal plain. **c**, Bivalve mollusc *Cuspidaria* sp. from the abyssal Weddell Sea and South African basins. **d**, Bivalve mollusc *Limatula* sp. from the Weddell Sea. **e**, Polychaete *Ophryotrocha* from the Weddell Sea. **f**, Nematode *Molgolaimus* sp. from the Weddell Sea. **h**, Isopod crustacean *Ischnomesus* sp. from the Weddell Sea. **g**, Isopod *Munnopsis* sp. from the western Weddell Sea.

amphipods and decapods but not for polychaetes, asteroids and ophiuroids<sup>12</sup>. The generally higher degree of eurybathy among Antarctic invertebrates was attributed to glacial–interglacial cycles of shelf ice advance and retreat, which periodically eliminated shelf faunas, pushing species into deeper water or causing their extinction. These cycles probably also led to regular pulses of migration in and out of Antarctica<sup>1</sup>. Experimental studies of the pressure and temperature tolerances of the pelagic larvae of the shallow-water Antarctic echinoid *Sterechinus neumayeri* suggested that embryonic stages potentially could penetrate into deep water<sup>13</sup>. The unusually deep shelf around Antarctica could also facilitate faunal exchanges between shelf and deep-sea habitats<sup>2</sup>.

ANDEEP data strongly suggest that a combination of emergence and submergence on Antarctic margins has led to an intermingling of species originating in shallow- and deep-water habitats (Fig. 3). The



**Figure 3 | Isopod phylogenetic relationships and multiple colonizations of the deep sea.** Bayesian consensus tree based on the complete 18S rRNA gene (excluding hypervariable regions; numbers on nodes represent statistical support) of 53 species of marine Asellota, and bathymetric distribution of the analysed species (broad lines). Family distribution (thin black lines) is based on ANDEEP data and the world list of isopods (www.nmnh.si.edu./iz/isopod/). Eye-bearing, shallow-water taxa are marked light blue; blind, deep-sea taxa are marked black. The family Acanthaspidiidae includes both shallow-water and deep-sea species (dark blue colour). The scattered distribution of the shallow-water forms within the tree, caused by the polyphyletic nature of the 'Janiridae', indicates a multiple colonization of the deep sea. In at least four cases, asellote families (including the Acanthaspidiidae) have invaded the deep sea. The Munnopidae are most speciose and include seven dominant deep-sea subfamilies. The bathymetric distribution of the analysed species provides evidence of polar submergence within the Stenetriidae, Joeropsididae and some janirid species, whereas polar emergence can be observed within the Acanthaspidiidae (genus *Acanthaspidia*), Dendrotionidae, Ischnomesidae, Desmosomatidae, Nannoniscidae and Munnopsidae.





These biogeographic patterns may be linked to larval ecology. The isopods, ostracods and nematodes have poor dispersal capabilities (isopods, for example, are brooders<sup>10</sup>) and hence reduced gene flow, making restricted species distributions more plausible, albeit difficult to prove. The recent discovery of cryptic isopod species in the Southern Ocean and Angola Basin support this assumption<sup>23</sup>. On the other hand, groups in which species are widely distributed, for example, Foraminifera, have easily transported propagules or larval stages<sup>24</sup>.

Our data can be used to test the suggested existence of global latitudinal diversity trends in the deep sea<sup>3,5,25,26</sup>. A decrease in isopod biodiversity with increasing latitude in the Southern Hemisphere has been claimed on the basis of data from up to 40° S (ref. 26), although species richness data point to considerable regional variation for isopods on continental slopes<sup>3,25–27</sup>. ANDEEP data, however, indicate high levels of diversity among certain animal groups in the deep Southern Ocean. Values of  $E(S_{100})$  (expected number of species per 100 individuals) for isopods from the Weddell and Scotia Seas (22–25 species) are comparable to those from many temperate and tropical slopes in the Southern Hemisphere<sup>25</sup>. Reported  $E(S_{100})$  and  $E(S_{200})$  values for bivalves, gastropods and isopods from the southeastern Weddell Sea (500–2,000 m) fall within the upper range of those from the tropics<sup>27</sup>. Isopod diversity is even higher ( $E(S_{100}) = 55$  species at 3,100 m) than on the southern Australian shelf (40 species)<sup>10</sup> and the Argentine basin<sup>28,29</sup>. Our results further indicate that these high levels of biodiversity extend to abyssal depths; for example,  $E(S_{100})$  for isopods is >30 at almost 5,000 m (Supplementary Fig. 4). Possibly, this is a regional phenomenon and diversity is lower in other parts of the deep Southern Ocean. Nevertheless, the extraordinary richness of species in ANDEEP samples when compared with literature data from tropical and temperate sites in the South Atlantic challenges the idea that deep-sea diversity is depressed towards the Antarctic continent. Moreover, the Atlantic is the youngest ocean. A longer period of faunal evolution and diversification in the Pacific and Indian Oceans may have led to greater species richness than in the Atlantic sector of the Southern Ocean. Continuing studies at the morphological and molecular levels of bathyal and abyssal Antarctic faunas may provide further insights into the evolutionary processes that have helped to shape these important communities.

## METHODS

A sediment profile imaging system (SPI) was used to obtain photographs and video footage of the seafloor and sediment profiles. Two Barnett-type, hydraulically damped multicorers equipped with 57-mm and 100-mm diameter tubes (25.5 and 78.6 cm<sup>2</sup> surface area respectively), and two box corers (2,500 cm<sup>2</sup> surface area) subdivided into 100 cm<sup>2</sup> 'vegematic' subcores recovered sediment samples with well-defined surface areas. Multicorers were used principally for the meiofauna (32–300 µm size range) and characterization of sediment parameters; box corers were used to sample the macrofauna (>300 µm). Two towed gears, an epibenthic sledge for macrofauna and an Agassiz trawl for megafauna, obtained material from a much larger but less well-defined area of seafloor and provided most of the taxonomic material in these two size classes. The sledge comprised a lower epibenthic net (500 µm mesh size) and an upper suprabenthic net (300 µm mesh size), each 1 m wide. It was pulled across the seabed for 10 min at a mean velocity of 1 knot. Total distances covered varied from 711 to 6,464 m. The 3-m-wide Agassiz trawl had a cod-end mesh size of 500 µm, except at stations 74, 78 and 81, at which a 10-mm mesh was used. It was fished in a similar way to the epibenthic sledge. Sledge catches were usually fairly clean, whereas the trawl often recovered large quantities of mud which required sieving. Specimens for morphological study were sorted either on board or in a shore laboratory. Metazoans for DNA extraction were picked from samples that had been transferred immediately after collection into pre-cooled 96% ethanol and kept for at least 48 h at –20 °C. Foraminifera for molecular study were picked from unfixed samples in an ice-chilled dish and either frozen in liquid nitrogen or the DNA individually isolated using guanidine lysis buffer.

肌裂解缓冲液

Received 13 September 2006; accepted 10 April 2007.

- Clarke, A. & Johnston, N. M. Antarctic marine benthic diversity. *Oceanogr. Mar. Biol.* **41**, 47–114 (2003).

- Clarke, A. in *Ecosystems of the World. Ecosystems of the Deep Oceans*, Vol. 28 (ed. Tyler, P. A.) 239–260 (Amsterdam, Elsevier, 2003).
- Gage, J. D. Diversity in deep-sea benthic Macrofauna: the importance of local ecology, the larger scale, history and the Antarctic. *Deep-sea Res. II* **51**, 1689–1708 (2004).
- Brandt, A. *et al.* The biodiversity of the deep Southern Ocean benthos. *Phil. Trans. Roy. Soc. B* **362**, 39–66 (2007).
- Rex, M. A., Crame, A., Stuart, C. T. & Clarke, A. Large-scale biogeographic patterns in marine mollusks: a confluence of history and productivity. *Ecology* **86**, 2288–2297 (2005).
- Sanders, H. L. & Hessler, R. R. Ecology of the deep-sea benthos. *Science* **163**, 1419–1424 (1969).
- McCann, K. S. The diversity–stability debate. *Nature* **405**, 228–233 (2000).
- Snelgrove, P. V. R. & Smith, C. R. A riot of species in an environmental calm: the paradox of the species-rich deep sea floor. *Oceanogr. Mar. Biol.* **40**, 311–342 (2002).
- Levin, L. A. *et al.* Environmental influences on regional deep-sea species diversity. *Annu. Rev. Ecol. Syst.* **32**, 51–93 (2001).
- Brandt, A. On the origin and evolution of Antarctic Peracarida (Crustacea, Malacostraca). *Sci. Mar.* **63**, 261–274 (1999).
- Hilbig, B. & Blake, J. A. Deep-sea polychaete communities in the northeast Pacific Ocean off the Gulf of the Farallones, California. *Bull. Mar. Sci.* **78**, 243–269 (2006).
- Brey, T. *et al.* Do Antarctic benthic invertebrates show an extended level of eurybathy? *Antarct. Sci.* **8**, 3–6 (1996).
- Tyler, P. A., Young, C. M. & Clarke, A. Temperature and pressure tolerances of embryos and larvae of the Antarctic sea urchin *Sterechinus neumayeri* (Echinodermata: Echinoidea): potential for deep-sea invasion from high latitudes. *Mar. Ecol. Prog. Ser.* **192**, 173–180 (2000).
- Pawlowski, J. *et al.* A note on the genetic similarity between shallow- and deep-water specimens of the Antarctic foraminifer *Epistominella vitrea*. *Deep-sea Res. II* (submitted).
- Cornelius, N. & Gooday, A. J. 'Live' (stained) deep-sea benthic foraminiferans in the western Weddell Sea: trends in abundance, diversity and taxonomic composition along a depth transect. *Deep-sea Res. II* **51**, 1571–1603 (2004).
- Murray, J. W. & Pudsey, C. J. Living (stained) and dead foraminifera from the newly ice-free Larsen Ice Shelf, Weddell Sea, Antarctica: ecology and taphonomy. *Mar. Micropaleontol.* **53**, 67–81 (2004).
- De Mesel, I., Lee, H. J., Vanhove, S., Vincx, M. & Vanreusel, A. Species diversity and distribution within the deep-sea nematode genus *Acantholaimus* on the continental shelf and slope in Antarctica. *Polar Biol.* **29**, 860–871 (2006).
- Hessler, R. R. & Thistle, D. On the place of origin of deep-sea isopods. *Mar. Biol.* **32**, 155–165 (1975).
- Kussakin, G. O. Peculiarities of the geographical and vertical distribution of marine isopods and the problem of deep sea fauna origin. *Mar. Biol.* **23**, 19–34 (1973).
- Raupach, M. J., Held, C. & Wägele, J.-W. Multiple colonization of the deep-sea by the Asellota (Crustacea: Peracarida: Asellota). *Deep-sea Res. II* **51**, 1787–1795 (2004).
- Vinogradova, N. G. Zoogeography of the abyssal and hadal zones. *Adv. Mar. Biol.* **32**, 325–387 (2004).
- Pawlowski, J. *et al.* Bipolar gene flow in deep-sea benthic foraminifera. *Molecular Ecol.* (submitted).
- Raupach, M. J., Malyutina, M., Brandt, A. & Wägele, J. W. Molecular data reveal a highly diverse species flock within the munnopsoid deep-sea isopod *Betamorphafusiformis* (Barnard, 1920) (Crustacea: Isopoda: Asellota) in the Southern Ocean. *Deep-sea Res. II* (submitted).
- Alve, E. & Goldstein, S. T. Propagule transport as a key method of dispersal in benthic foraminifera (Protista). *Limnol. Oceanogr.* **48**, 2163–2170 (2003).
- Poore, G. C. B. & Wilson, G. D. F. Marine species richness. *Nature* **361**, 597–598 (1993).
- Rex, M. A., Stuart, C. T., Hessler, R. R., Allen, J. A., Sanders, H. L., & Wilson, G. D. F. Global-scale latitudinal patterns of species diversity in the deep-sea benthos. *Nature* **365**, 636–639 (1993).
- Brey, T. *et al.* Antarctic benthic diversity. *Nature* **368**, 297 (1994).
- Hessler, R. R. & Wilson, G. D. F. in *Evolution, Time and Space: The Emergence of the Biosphere*. (eds Sims, R.W., Price, J.H. & Whalley, P.E.S.) 227–254 (Academic Press, London, 1983).
- Brandt, A. Zur Besiedlungsgeschichte des antarktischen Schelfes am Beispiel der Isopoda (Crustacea, Malacostraca). *Ber. Polarforsch.* **98**, 1–240 (1991).

Supplementary Information is linked to the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

**Acknowledgements** Financial support for the ANDEEP I–III expeditions was provided by the German Science Foundation. We are also grateful for support from the Swiss National Science Foundation and UK Natural Environment Research Council. S.N.B. received financial support from the DAAD, CAPES, Scar and Synthesys. We thank L. Excoffier, J. Guiard, J. Guiard, D. Longet and B. Lecroq for help in obtaining the foraminiferal sequence data. We are grateful to D. Fütterer, chief scientist on *Polarstern* cruise ANT XIX/3–4, and E. Fahrbach, chief scientist on *Polarstern* cruise ANT XXII/3, and to the captain and crew of RV *Polarstern*, for help

on board. The DZMB is thanked for financial support of sorting animals from samples and for a research grant for F. Kavanagh, who kindly identified the ANDEEP III Ischnomesidae (Isopoda) and compared these with those species from ANDEEP I and II. We are grateful to many colleagues for constant help on board, helpful and stimulating discussions, and comments, which improved the manuscript. This is ANDEEP publication no. 85 and CAML publication no. 6. This publication also contributes to the CoML field project CeDAMar.

**Author Contributions** A.B. and A.J.G. assembled and wrote most of the text and were responsible for the theory. J.H. contributed to sedimentology and B. Diaz to sediment analysis, bioturbation and megafauna visible in underwater images. B.

Danis and D.C.G. worked on archaeobacteria. Foraminiferal data are from N.C., T.C., A.J.G. and J.P.; J.P. was responsible for the molecular work on the Foraminifera. I.D.M. and A.V. delivered the data for Nematoda. A.B., S.B., W.B., M.C., S.K. and M.M. were responsible for the isopod analysis, M.R. was mainly responsible for the molecular isopod data. S.N.B. was responsible for the Ostracoda, B.E. for the Polychaetes, K.L. for the Mollusca and D.J. for the Porifera analysis.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to A.B. ([abrandt@zoologie.uni-hamburg.de](mailto:abrandt@zoologie.uni-hamburg.de)).