

Draft Genome Sequences of Two *Legionella dumoffii* Strains, TEX-KL and NY-23

Tian Qin, Yujun Cui, Zhong Cen, Ting Liang, Hongyu Ren, Xianwei Yang, Xiangna Zhao, Zhangyi Liu, Li Xu, Dongfang Li, Youtao Song, Ruifu Yang, Zhujun Shao and Yajun Song
J. Bacteriol. 2012, 194(5):1251. DOI: 10.1128/JB.06352-11.

Updated information and services can be found at:
<http://jb.asm.org/content/194/5/1251>

	<i>These include:</i>
REFERENCES	This article cites 16 articles, 7 of which can be accessed free at: http://jb.asm.org/content/194/5/1251#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://jb.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Draft Genome Sequences of Two *Legionella dumoffii* Strains, TEX-KL and NY-23

Tian Qin,^a Yujun Cui,^b Zhong Cen,^c Ting Liang,^{b,d} Hongyu Ren,^a Xianwei Yang,^c Xiangna Zhao,^b Zhangyi Liu,^c Li Xu,^{b,e} Dongfang Li,^c Youtao Song,^d Ruifu Yang,^b Zhujun Shao,^a and Yajun Song^b

State Key Laboratory for Infectious Disease Prevention and Control and National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, People's Republic of China^a; State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing, People's Republic of China^b; BGI-Shenzhen, Shenzhen, People's Republic of China^c; Department of Microbiology, Liaoning University, Shenyang, People's Republic of China^d; and The Second Affiliated Hospital of Hebei Medical University, Shijiazhuang, People's Republic of China^e

***Legionella (Fluoribacter) dumoffii* is one of the agents causing Legionnaires' disease. Here, we used Illumina second-generation sequencing technology to decipher for the first time the whole-genome sequences of two strains of this species, TEX-KL and NY-23. The assembly results for both strains consist of one chromosome and two plasmids.**

Legionella dumoffii, a pathogen that is a less common cause of Legionnaires' disease than *Legionella pneumophila* (6), was first isolated in 1979 as atypical *Legionella*-like organisms (ALLO) (4, 9), and it was designated as a new species in 1980 (1). *L. dumoffii* lives in soil and freshwater environments, and it is capable of surviving in human alveolar macrophages and epithelial cells as well (10). Infection by *L. dumoffii* can cause human pneumonia and accidentally induce other diseases, such as prosthetic-valve endocarditis and septic arthritis (7, 15). Based on its phenotypic differences from other species of the *Legionella* genus and the experimental results of DNA hybridization, Brown et al. suggested that *L. dumoffii* be transferred to the genus *Fluoribacter* as *F. dumoffii*, which is not yet widely accepted (2).

According to the Genomes Online Database (<http://www.genomesonline.org>), there is no genome sequence for *L. dumoffii* reported so far, and only one plasmid sequence from this species was described by us (11). Here, we present the whole-genome sequences of two *L. dumoffii* strains, TEX-KL (ATCC 33343) and NY-23 (ATCC 33279). TEX-KL was initially isolated from a post-mortem lung specimen of a patient with fatal atypical pneumonia in Houston, TX (9), and NY-23 was isolated from a cooling tower in New York (4).

The strains were cultured on buffered charcoal-yeast extract (BCYE) agar, and DNAs were extracted as previously described (12). Whole-genome sequencing was performed using Illumina HiSeq 2000 (Illumina, Inc., United States) by generating paired-end libraries (500 bp and 6 kb) following the manufacturer's instructions. The read lengths were 90 bp and 49 bp for each library, from which 400 Mb and 210 Mb, respectively, of high-quality data were generated. The paired-end reads were first *de novo* assembled using SOAPdenovo version 1.06 (13), and then the scaffolds were manually connected according to 6-kb paired-end relationships. The coding sequences (CDS) were predicted by using Glimmer version 3.02 (5), and homologous comparison to the NCBI nonredundant public database for function annotations was performed by BLAST. The rRNA and tRNA were identified using RNAmmer (8) and tRNAscan-SE 1.21 (14), respectively, and the insertion sequence (IS) elements were annotated with ISSaga (16).

We obtained three scaffolds for each strain, which were identified as one circular chromosome and two plasmid sequences for each of them. The G+C contents of the chromosome are 39.7%,

and the G+C contents of the plasmids are slightly lower (37.5 to 39.5%). The average nucleotide identity (ANI) between the chromosomes of the two strains is 99.4%. For strain TEX-KL, the smaller plasmid (pLD-TEX-KL, 66.5 kb) is identical to the one published previously (11). The smaller plasmid in strain NY-23 (pLDNY1, 81.8 kb) is similar to pLD-TEX-KL, with 51.3-kb common sequences identified between them (identity, >80%). The larger plasmids, the 126.5-kb pLDTK for TEX-KL and the 138.3-kb pLDNY2 for NY-23, were found to be homologous with pLPP, a previously reported plasmid of *L. pneumophila* strain Paris (132 kb) (3). There are 3,562 predicted CDS in TEX-KL (3,370 in chromosome and 192 in the plasmids), while there are 3,637 CDS in NY-23 (3,409 in the chromosome and 228 in the plasmids).

Nucleotide sequence accession numbers. This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under accession numbers AGVT00000000 (Tex-KL) and AGVU00000000 (NY 23). The versions described in this paper are the first versions, AGVT01000000 and AGVU01000000.

ACKNOWLEDGMENTS

This work was supported by the National Key Program for Infectious Diseases of China (grants 2008ZX10004-001-C and 2012ZX10004-216), the State Key Development Program for Basic Research of China (grant 2009CB522600), and the Science Foundation for the State Key Laboratory for Infectious Disease Prevention and Control from China, Chinese CDC (grant 2011SKLID202).

REFERENCES

- Brenner D, et al. 1980. *Legionella bozemanii* sp. nov. and *Legionella dumoffii* sp. nov.: classification of two additional species of *Legionella* associated with human pneumonia. *Curr. Microbiol.* 4:111–116.
- Brown A, Garrity GM, Vichers RM. 1981. *Fluoribacter dumoffii* (Brenner

Received 2 December 2011 Accepted 12 December 2011

Address correspondence to Yajun Song, songyajun88@yahoo.com.cn, or Zhujun Shao, shaozhujun@icdc.cn.

Tian Qin, Yujun Cui, and Zhong Cen contributed equally to this work.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.06352-11

- et al.) comb. nov. and *Fluoribacter gormanii* (Morris et al.) comb. nov. Int. J. Syst. Bacteriol. 31:111–115.
3. Cazalet C, et al. 2004. Evidence in the *Legionella pneumophila* genome for exploitation of host cell functions and high genome plasticity. Nat. Genet. 36:1165–1173.
 4. Cordes LG, Wilkinson HW, Gorman GW, Fikes BJ, Fraser DW. 1979. Atypical *Legionella*-like organisms: fastidious water-associated bacteria pathogenic for man. Lancet ii:927–930.
 5. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. Bioinformatics 23:673–679.
 6. Fang GD, Yu VL, Vickers RM. 1989. Disease due to the *Legionellaceae* (other than *Legionella pneumophila*). Historical, microbiological, clinical, and epidemiological review. Medicine (Baltimore) 68:116–132.
 7. Flendrie M, et al. 2011. Septic arthritis caused by *Legionella dumoffii* in a patient with systemic lupus erythematosus-like disease. J. Clin. Microbiol. 49:746–749.
 8. Lagesen K, et al. 2007. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids Res. 35:3100–3108.
 9. Lewallen KR, et al. 1979. A newly identified bacterium phenotypically resembling, but genetically distinct from, *Legionella pneumophila*: an isolate in a case of pneumonia. Ann. Intern. Med. 91:831–834.
 10. Maruta K, et al. 1998. Entry and intracellular growth of *Legionella dumoffii* in alveolar epithelial cells. Am. J. Respir. Crit. Care Med. 157:1967–1974.
 11. Qin T, et al. 2007. Complete nucleotide sequence of pLD-TEX-KL, a 66-kb plasmid of *Legionella dumoffii* TEX-KL strain. Plasmid 58:261–268.
 12. Qin T, et al. 2009. Conjugative plasmid pLD-TEX-KL promotes growth of host bacterium *Legionella dumoffii* at low temperatures. Arch. Microbiol. 191:543–551.
 13. Rohde H, et al. 2011. Open-source genomic analysis of Shiga-toxin-producing *E. coli* O104:H4. N. Engl. J. Med. 365:718–724.
 14. Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. Nucleic Acids Res. 33:W686–W689.
 15. Tompkins LS, Roessler BJ, Redd SC, Markowitz LE, Cohen ML. 1988. *Legionella* prosthetic-valve endocarditis. N. Engl. J. Med. 318:530–535.
 16. Varani AM, Siguier P, Gourbeyre E, Charneau V, Chandler M. 2011. ISSaga is an ensemble of web-based methods for high throughput identification and semi-automatic annotation of insertion sequences in prokaryotic genomes. Genome Biol. 12:R30.