

# Bacterial Quorum-Sensing Network Architectures

Wai-Leung Ng<sup>1</sup> and Bonnie L. Bassler<sup>1,2</sup>

<sup>1</sup>Department of Molecular Biology, Princeton University, Princeton, New Jersey 08544-1014; email: bbassler@princeton.edu

<sup>2</sup>Howard Hughes Medical Institute, Chevy Chase, Maryland 20815-6789

Annu. Rev. Genet. 2009. 43:197–222

First published online as a Review in Advance on August 17, 2009

The *Annual Review of Genetics* is online at [genet.annualreviews.org](http://genet.annualreviews.org)

This article's doi:  
10.1146/annurev-genet-102108-134304

Copyright © 2009 by Annual Reviews.  
All rights reserved.

0066-4197/09/1201-0197\$20.00

## Key Words

autoinducers, signal transduction, feedback regulation, small RNAs

## Abstract

Quorum sensing is a cell-cell communication process in which bacteria use the production and detection of extracellular chemicals called autoinducers to monitor cell population density. Quorum sensing allows bacteria to synchronize the gene expression of the group, and thus act in unison. Here, we review the mechanisms involved in quorum sensing with a focus on the *Vibrio harveyi* and *Vibrio cholerae* quorum-sensing systems. We discuss the differences between these two quorum-sensing systems and the differences between them and other paradigmatic bacterial signal transduction systems. We argue that the *Vibrio* quorum-sensing systems are optimally designed to precisely translate extracellular autoinducer information into internal changes in gene expression. We describe how studies of the *V. harveyi* and *V. cholerae* quorum-sensing systems have revealed some of the fundamental mechanisms underpinning the evolution of collective behaviors.

**Quorum sensing:**

a cell-cell communication process used by bacteria to coordinate gene expression in response to changes in population density

**LCD:**

low-cell-density

**HCD:**

high-cell-density

**Autoinducers:**

extracellular signal molecules produced and detected by bacteria to monitor cell population density

**AHL:**

acyl homoserine lactone autoinducer

## QUORUM SENSING

Bacterial processes such as biofilm formation, virulence factor secretion, bioluminescence, antibiotic production, sporulation, and competence for DNA uptake are often critical for survival. However, these behaviors are seemingly futile if performed by a single bacterium acting alone. Yet, we know that bacteria perform these tasks effectively. How do bacteria manage? We now understand that, through a process called quorum sensing, bacteria synchronously control gene expression in response to changes in cell density and species complexity. Quorum sensing allows bacteria to switch between two distinct gene expression programs: one that is favored at low-cell-density (LCD) for individual, asocial behaviors, and another that is favored at high-cell-density (HCD) for social, group behaviors (reviewed in 87, 88, 133, 138).

The fundamental steps involved in detecting and responding to fluctuations in cell number are analogous in all known quorum-sensing systems. First, low molecular weight molecules called autoinducers are synthesized intracellularly. Second, these molecules are either passively released or actively secreted outside of the cells. As the number of cells in a population increases, the extracellular concentration of autoinducer likewise increases. Third, when autoinducers accumulate above the minimal threshold level required for detection, cognate receptors bind the autoinducers and trigger signal transduction cascades that result in population-wide changes in gene expression. Thus, quorum sensing enables cells in a population to function in unison and in so doing, they carry out behaviors as a collective.

In this review, following a short synopsis of canonical quorum-sensing systems, we focus on the quorum-sensing systems of *Vibrio harveyi* and *Vibrio cholerae*. Studies of these







two paradigmatic systems have revealed fundamental molecular mechanisms underpinning small-molecule biosynthesis, signal detection and transduction, information processing, small RNA-mediated post-transcriptional control of mRNA levels, and transcriptional gene regulation (37, 38, 57, 75, 97, 124, 134). Furthermore, analysis of the similarities and differences in these two related quorum-sensing systems, as well as assessing differences between these two systems and other bacterial signal transduction systems, provides insight into how evolution has incorporated distinct features into bacterial sensory networks that solve unique biological information processing challenges.

## LUXIR-TYPE QUORUM-SENSING SYSTEMS IN GRAM-NEGATIVE BACTERIA

Acyl homoserine lactones (AHLs) are a major class of autoinducer signal used by Gram-negative proteobacteria for intraspecies quorum sensing. AHLs are composed of homoserine lactone (HSL) rings carrying acyl chains of C<sub>4</sub> to C<sub>18</sub> in length (25). These side chains harbor occasional modification, notably at the C<sub>3</sub> position, or unsaturated double bonds (**Figure 1a**). The first AHL autoinducer and its cognate regulatory circuit was discovered in the bioluminescent marine bacterium *Vibrio fischeri*, which colonizes the light organ of the Hawaiian Bobtail Squid *Euprymna scolopes* (100). The nutritious environment inside the light organ of the squid allows the bacteria to grow to high cell density, and using quorum sensing, to activate expression of the luciferase operon. The squid host uses the bacterial-produced light to counter-illuminate itself in an antipredation strategy (100). Two proteins, LuxI and LuxR, are essential for quorum-sensing control of

### Figure 1








Structures of bacterial autoinducers. (a) Homoserine lactone autoinducers produced by different Gram-negative bacteria. (b) Amino acid sequences of three peptide autoinducers, ComX, CSF, and CSP, produced by Gram-positive bacteria. The underlined tryptophan in *Bacillus subtilis* ComX is isoprenylated. The four different AIPs produced by *S. aureus*. (c) DPD, the precursor to AI-2. In the presence of boron, AI-2 exists as S-THMF-borate. In the absence of boron, AI-2 exists as R-THMF. (d) Structure of *V. cholerae* CAI-1 and Amino-CAI-1. (e) Structure of the PQS autoinducer of *Pseudomonas aeruginosa*.


**a****R group****V. fischeri (LuxI)****V. harveyi (LuxM)****P. aeruginosa (RhII)****P. aeruginosa (LasI)****c****S-TMHF-borate  
(Vibrios)****R-TMHF  
(Salmonella)****b**ADPITRQWGD**B. subtilis (ComX)**

ERGMT

**B. subtilis (CSF)**

EMRLSKFFRDFILQRKK

**S. pneumoniae (CSP)****S. aureus (AgrD)****AIP-I****AIP-III****AIP-II****AIP-IV****d****e**



**Figure 2**

A canonical Gram-negative LuxIR-type quorum-sensing system. Red pentagons denote AHL autoinducers.

bioluminescence in *V. fischeri* (Figure 2). LuxI is the synthase of the quorum-sensing autoinducer *N*-3-(oxo-hexanoyl)-homoserine lactone (3OC6HSL, Figure 1) (19, 104). LuxI catalyzes acylation and lactonization reactions between the substrates *S*-adenosylmethionine (SAM) and hexanoyl-ACP (81, 104). Following synthesis, 3OC6HSL diffuses freely in and out of the cell, and its concentration increases as the cell density of the population increases (48). LuxR is the cytoplasmic receptor for 3OC6HSL as well as the transcriptional activator of the luciferase *luxICDABE* operon (18, 19). Without the 3OC6HSL ligand, the LuxR protein is unstable and is rapidly degraded. When 3OC6HSL accumulates, it is bound by LuxR, and the LuxR-AHL complex recognizes a consensus binding sequence (lux box) upstream of the *luxICDABE* operon and activates its expression (114). The *luxAB* genes encode the subunits of luciferase, and *luxCDE* encode the fatty acid reductase complex, which produces and recycles the luciferase aldehyde substrate (74). Because expression of *luxI* is also activated by 3OC6HSL-bound LuxR, when the quorum-sensing circuit engages, autoinducer production is induced, and the surrounding environment is flooded with the signal molecule. This autoinduction positive feedback loop is presumed to enforce synchrony as the population of cells switches from LCD mode to HCD quorum-sensing mode.

**SAM:**  
S-adenosylmethionine  
**ACP:** acyl carrier  
protein

The LuxI/LuxR regulatory system of *V. fischeri* is considered the paradigm for the control of gene expression by quorum sensing in Gram-negative bacteria (Figure 2) (19). Homologs of *luxI* and *luxR* have been identified in a large number of bacterial genomes, and these other LuxIR-type quorum-sensing systems control global cell-density-dependent gene expression (11). Positive feedback loops consisting of LuxR-type proteins activating *luxI*-type gene expression are commonly wired into these AHL quorum-sensing systems (18, 26, 108). Some well studied AHL quorum-sensing systems include the LasI/LasR-RhlI/RhlR system of *Pseudomonas aeruginosa* that controls virulence factor gene expression and biofilm formation (89–91, 139), the TraI/TraR system of *Agrobacterium tumefaciens* that regulates transfer of the oncogenic Ti plasmid to the plant host (26, 40, 96), and the EsaI/EsaR system in *Pantoea stewartii* that controls exopolysaccharide production, adhesion, and plant colonization (130, 131). In an interesting twist on these circuits, the EsaI/EsaR system functions reciprocally to other LuxIR-type systems. Unliganded EsaR binds DNA and represses transcription. Autoinducer binding to EsaR promotes DNA release, and target gene expression (79, 129).

AHL autoinducer molecules are typically unique in that a particular AHL molecule is detected only by the species that produces it. Therefore, it is suggested that AHL-type quorum-sensing systems predominately foster intraspecies cell-cell communication. Signal specificity is attributed to the fact that LuxR-type protein folding requires the presence of the AHL ligand (107, 126, 148, 149). For example, in the absence of its cognate ligand (3OC8HSL), TraR is unstructured and rapidly degraded (148, 149). Supporting this notion is the finding that the crystal structure of the 3OC8HSL-TraR complex reveals that the ligand is completely buried within the protein (127, 145). Solution structure of the ligand binding domain complexed with C8HSL of SdiA, a TraR homolog in *Escherichia coli*, reveals similar organization (144). Although *P. aeruginosa* LasR shares low overall homology

with TraR and SdiA, the LasR ligand binding domain nonetheless shows significant structural similarity to these two proteins (8). Importantly, the putative signal binding pocket of LasR is more spacious than that of TraR and SdiA. The presumption is that this arrangement accommodates the larger cognate ligand, 3OC12HSL (8).

Biochemical analyses of LuxR-type proteins suggest that there exist three classes of these receptors (reviewed in 106). Class 1 receptors (e.g., LasR) require AHL for folding and have exquisitely tight affinity for their ligands. Class 2 receptors (e.g., LuxR of *V. fischeri*) also require AHLs for folding, but the mature proteins bind to their ligands reversibly. Class 3 receptors (e.g., EsaR) do not require AHLs for folding and have reversible ligand binding. The absolute requirement for AHLs for protein folding in Class 1 and 2 LuxR-type receptors suggests that these receptors are refractory to sudden increases in AHL concentrations in the environment because protein translation is a relatively slow process (106). It is further suggested that the difference in ligand binding affinity between these receptors is the main criterion controlling their robustness to perturbation (106). For example, the folded Class 1 receptor TraR remains active for a prolonged period of time even when its cognate AHL is removed (63), whereas the folded Class 2 receptor LuxR is inactivated within minutes after its cognate AHL is removed (48).

Two different structures of AHL synthases, LasI from *P. aeruginosa* that synthesizes 3OC12HSL and EsaI from *P. stewartii* that synthesizes 3OC6HSL, have been reported (31, 136). By comparing the two structures, it appears that although LasI and EsaI synthesize different molecules, the core domains that contain 74 conserved amino acid residues are highly similar (31, 136). However, the acyl chain binding pockets differ dramatically. The binding pocket of EsaI sits in an enclosed cavity surrounded by numerous other residues (136). By contrast, the substrate binding pocket in LasI is an elongated tunnel (31). These structural features suggest that EsaI can

only accommodate substrates with relatively short acyl-chain lengths, whereas LasI has no steric restriction on the substrate acyl-chain length (31, 136). Thus, it is not understood how LasI selects only the appropriate substrate (i.e., 3OC<sub>12</sub>-ACP) for reaction. One hypothesis is that substrates with acyl chains longer than C<sub>12</sub> are used by the cell for other essential processes such as LPS biosynthesis and this limits their availability to LasI (31). As with LasI and EsaI, many other LuxI-type AHL quorum-sensing synthases use the intracellular fatty acid pool as the source of substrate for AHL synthesis. One notable exception is the newly discovered RpaI/RpaR system in the photosynthetic bacterium *Rhodospseudomonas palustris*. In this case, the signal, *p*-coumaroyl-HSL, is generated from *p*-coumaric acid obtained from the extracellular environment (103). Given that *p*-coumarate is a major by-product of lignin degradation in plants, it is proposed that the signal *p*-coumaroyl-HSL is used for both intraspecies signaling among bacterial cells and interkingdom signaling between bacterial and plant cells (103).


## TWO-COMPONENT QUORUM-SENSING SYSTEMS IN GRAM-POSITIVE BACTERIA

Gram-positive bacteria primarily use modified oligopeptides as autoinducers in quorum-sensing controlled gene expression systems (**Figure 1b** and **Figure 3**) (35, 45, 110). Because peptides are impermeable to biological membranes, secretion of quorum-sensing peptides is usually mediated by specialized transporters. In addition, modifications to the initially synthesized peptides, such as processing and cyclization, are often associated with secretion (**Figure 1**) (35, 44, 69, 70, 110). One major difference between LuxIR-based and oligopeptide-based quorum-sensing systems is the location of the cognate receptors; whereas the LuxR-type receptors are cytoplasmic, the sensors for oligopeptide autoinducers in Gram-positive bacteria are membrane-bound.

---

**LPS:**  
lipopolysaccharide

---



**Figure 3**

A canonical Gram-positive two-component-type quorum-sensing system. Blue octagons denote processed/modified peptide autoinducers.

The membrane-bound receptors, so-called two-component signaling proteins, transduce information via a series of phosphorylation events (39, 41, 109).

A typical two-component system consists of a membrane-bound histidine kinase receptor and a cognate cytoplasmic response regulator, which functions as a transcriptional regulator (39, 41, 109). A general model for oligopeptide-mediated quorum sensing is depicted in **Figure 3**. As in AHL quorum-sensing systems, the concentration of secreted oligopeptide autoinducer increases as the cell density increases. Peptide binding to the membrane-bound histidine kinase receptor stimulates its intrinsic autophosphorylation activity, resulting in ATP-driven phosphorylation of a conserved histidine residue (H) in the cytoplasm. The phosphate group is subsequently transferred to the conserved aspartate residue (D) of a cognate response regulator. Phosphorylated response regulators are active and they function as DNA-binding transcription factors to modulate expression of target genes. In many cases, the genes encoding the oligopeptide autoinducer precursor, the histidine kinase receptor, and the response regulator form an operon, and its expression is auto-induced by quorum sens-

ing (45, 95). In such cases, similar to what was described above for the LuxR/LuxI systems, this configuration produces positive feedback and accelerates the transition from the LCD to the HCD quorum-sensing mode of gene expression. Examples of peptide-based quorum-sensing systems include the ComD/ComE system of *Streptococcus pneumoniae* that controls competence development (94), the AgrC/AgrA system of *Staphylococcus aureus* that controls pathogenesis (45), and the ComP/ComA system of *Bacillus subtilis* that controls competence and sporulation (69).

Unlike Gram-negative bacterial AHLs, Gram-positive peptide autoinducers are not variations on a single core molecule. Rather, peptide autoinducers are genetically encoded, and thus each species of bacteria is capable of producing a peptide signal with a unique sequence (**Figure 1b**). Consistent with this, although Gram-positive quorum-sensing receptors are members of the histidine kinase protein family and thus share overall homology, little homology exists in their transmembrane ligand sensing domains and this likely determines their specificity (69, 87, 94). Although no Gram-positive quorum-sensing receptor has yet been crystallized, elegant genetic and biochemical studies have defined the *S. aureus* Agr quorum-sensing receptor-ligand interaction, and it stands as the paradigm for understanding signal transduction in peptide quorum-sensing systems (Reviewed in 87).

The *S. aureus* Agr autoinducer is denoted AIP (for Auto-inducing peptide) and is encoded as a longer precursor peptide by the *agrD* gene. Processing (truncation and cyclization) and secretion occurs via the AgrB transporter. Extracellular AIP is detected by the AgrC histidine kinase receptor and signal transduction occurs by phosphorelay to the AgrA response regulator (87). There are four *S. aureus* AIP specificity groups (I-IV) that are defined by the particular AIP peptide sequence (**Figure 1b**). The mature AIPs are seven to nine residues in length with a five-membered ring formed between the sulfur atom from a central cysteine and the C-terminus via a thiolactone bond. The

two bulky hydrophobic residues of each AIP are involved in AgrC binding, and the ring structure is critical for activity (67, 70, 141). The AIPs and their cognate receptors are coevolving such that productive signaling interactions occur only between a particular AIP and the hypervariable region of its cognate AgrC transmembrane domain (42, 44). Remarkably, interaction between an AIP and a noncognate AgrC receptor inhibits quorum sensing (42, 44, 66). Mutagenesis studies pinpoint the residues in the extracellular loop connecting transmembrane helices 3 and 4 as critical for AIP discrimination, whereas mutations in Ile171 broaden AIP specificity (27, 28, 43).

### ADDITIONAL FEATURES OF GRAM-NEGATIVE AND GRAM-POSITIVE QUORUM-SENSING SYSTEMS

Integration of additional regulatory features is common in both Gram-negative and Gram-positive quorum-sensing systems. We name only a few here. In *P. aeruginosa*, the CRP homolog Vfr is activated by binding the second messenger molecule cyclic-AMP (140). Activated Vfr induces transcription of *lasR* when *P. aeruginosa* enters into stationary phase (1). One of the target genes activated by the *S. pneumoniae* ComD/ComE system is *comX*, which encodes an alternative sigma factor that is essential for transcription of a set of late competence genes encoding proteins involved in genetic exchange such as DNA uptake and recombination (55, 61, 62). Interestingly, competence occurs for only a short period of time because ComX disappears soon after competence has developed. The ATP-dependent protease ClpEP has been implicated in specific degradation of ComX and termination of competence (116). These and other accessory control mechanisms allow bacteria to integrate environmental cues in addition to autoinducer information into their quorum-sensing networks presumably to extract maximal information from their surroundings and fine tune their transitions into and out of LCD and HCD gene expression programs.

### THE QUORUM-SENSING NETWORK ARCHITECTURE IN *V. harveyi* AND *V. cholerae*

Although LuxIR systems similar to that of *V. fischeri* have been identified in distantly related bacteria, analyses of quorum sensing in other *Vibrios*, most notably *V. harveyi* and *V. cholerae*, show that their systems do not conform to the common LuxIR theme. These two species possess neither *luxI* nor *luxR* genes similar to *V. fischeri* and other Gram-negative quorum-sensing bacteria. However, *V. harveyi* and *V. cholerae* do possess sets of quorum-sensing components that are highly similar to one another (37, 57, 75, 124, 147). This close similarity initially suggested functional equivalence among the components, however, detailed analyses of the roles of the individual components and the overall function of the networks reveal that these two quorum-sensing circuits operate by surprisingly different means (37, 57, 75, 125). We hypothesize that these differences have evolved to allow these two closely related species to adapt to their drastically different lifestyles.

*V. harveyi* is a free living marine bacterium and is an important pathogen of marine organisms (2). *V. cholerae*, by contrast, is the etiological agent of the disease cholera, and its life cycle consists of alternations between human hosts and the aquatic environment (20). Quorum sensing in *V. harveyi* activates bioluminescence and metalloprotease production and represses type III secretion (9, 36). In *V. cholerae*, quorum sensing represses biofilm formation and virulence factor production and activates protease production (32, 46, 146, 147). Quorum sensing also promotes genetic exchange between *V. cholerae* cells in the presence of chitin, which is believed to be important for serogroup conversion (6, 7, 73).

We first outline the quorum-sensing circuit in *V. harveyi*, a hybrid of the canonical Gram-negative and Gram-positive bacterial quorum-sensing systems (**Figure 4**). As in other Gram-negative quorum-sensing bacteria, *V. harveyi* produces, detects, and responds to an AHL autoinducer (3OHC4-HSL)

---

**HAI-1:** *V. harveyi*  
autoinducer 1

**AI-2:** autoinducer 2

**CAI-1:** *V. cholerae*  
autoinducer 1

**DPD:** 4,5-dihydroxy-  
2,3-pentanedione

**sRNAs:** small RNAs

---

denoted HAI-1 for *V. harveyi* autoinducer 1 (**Figure 1a**) (10). Two additional autoinducer molecules called AI-2 and CAI-1 are produced and detected by *V. harveyi* (**Figure 1c** and **d**, respectively). AI-2 is a set of interconverting molecules derived from the shared precursor 4,5-dihydroxy-2,3-pentanedione (DPD) (105). The active form of AI-2 in *V. harveyi* contains boron (13). CAI-1 has not been purified from *V. harveyi*, but it is presumed to be related to CAI-1 produced by *V. cholerae*, which has been identified as (S)-3-hydroxytridecan-4-one (38).

HAI-1 is synthesized by the LuxM synthase, which shows no homology to the LuxI-type AHL synthases although it carries out the same biochemistry (3, 4). DPD is synthesized by the LuxS enzyme, and *luxS* exists in hundreds of bacterial genomes (105, 117). We provide further detail about LuxS and AI-2 below. CAI-1 is synthesized by the CqsA synthase, and *cqsA* shows sequence homology to aminotransferases. CqsA homologs have been identified in several sequenced *Vibrio* genomes as well as *Legionella pneumophila* (37, 38, 75, 113, 122, 123).


Detection of the *V. harveyi* autoinducers does not occur through LuxR-type proteins. Rather, membrane-bound histidine kinases act as cognate receptors for all three autoinducers (**Figure 4**). HAI-1 is detected by the LuxN histidine kinase (24, 37, 120). AI-2 is detected by the periplasmic protein LuxP in complex with the LuxQ histidine kinase (4, 37, 84, 85). CAI-1 is detected by the CqsS histidine kinase (37, 38, 75). LuxN, LuxQ, and CqsS are bi-functional two-component enzymes that possess both kinase and phosphatase activities. At LCD (**Figure 4a**), the receptors are devoid of their respective ligands, and in this mode, their kinase activities predominate, resulting in phosphorylation of conserved histidine residues by ATP. The phosphate group is next transferred to the conserved aspartate residue located in the receiver domain at the C-terminus of each receptor. Phosphate from all three receptors is subsequently transduced to a single phosphotransfer protein, LuxU. LuxU transfers the phosphate to a response regulator called LuxO. LuxO belongs to the NtrC family of

response regulators and requires phosphorylation to act as a transcriptional activator (5, 23, 58).

Together with  $\sigma^{54}$ -loaded RNA polymerase, phosphorylated LuxO (LuxO-P) activates transcription of genes encoding five small regulatory RNAs (sRNAs) called Qrr1–5 (**Figure 4a**). The main target of the Qrr sRNAs is the mRNA encoding the quorum-sensing master transcriptional regulator LuxR. *V. harveyi* LuxR is not similar to the LuxRs described above in canonical LuxIR-type quorum-sensing systems. At LCD, the Qrr sRNAs are transcribed, and with the assistance of the RNA chaperone Hfq, these sRNAs base pair with and destabilize the *luxR* mRNA transcript (124). Therefore, at LCD, LuxR protein is not made. When autoinducer concentration is above the threshold level required for detection (e.g., at HCD, **Figure 4b**), binding of autoinducers to the cognate receptors switches the receptors from kinases to phosphatases. Phosphate flow in the signal transduction pathway is reversed, resulting in dephosphorylation and inactivation of LuxO. Therefore, at HCD, *qrr1–5* are not transcribed, *luxR* mRNA is stabilized, and LuxR protein is produced. LuxR acts as both a transcriptional activator and a transcriptional repressor. In addition to the luciferase operon, LuxR regulates at least another 50 genes including those encoding the type III secretion apparatus and metalloproteases (36, 97, 134).


The components and the wiring of the *V. cholerae* quorum-sensing system (**Figure 5**) appear extremely similar to their *V. harveyi* counterparts with only two notable differences. First, *V. cholerae* does not have the LuxM synthase and does not make HAI-1. Consistent with this, *V. cholerae* also does not have the LuxN sensor, and it does not detect HAI-1. Second, only four sRNAs genes lie downstream of LuxO-P in the *V. cholerae* cascade (57). The functional homolog of the *V. harveyi* LuxR master regulator is called HapR in *V. cholerae*. HapR, analogous to LuxR, acts as both an activator and a repressor of gene expression. At HCD (**Figure 5b**), HapR activates a gene encoding the Hap protease (46) and represses





**Figure 4**

The *V. harveyi* quorum-sensing circuit. (a) Signal transduction at LCD. During this stage, autoinducer levels are low and the LuxN, LuxPQ, and CqsS receptors function as kinases. LuxO is phosphorylated, the Qrr1–5 sRNAs are transcribed, and LuxR protein is not produced. (b) Signal transduction at HCD. During this stage, autoinducer levels are high and the LuxN, LuxPQ, and CqsS receptors function as phosphatases. LuxO is unphosphorylated, Qrr1–5 sRNAs are not transcribed, and LuxR protein is produced. Solid and dotted lines denote regulatory factors that are produced and not produced, respectively.



**Figure 5**

The *V. cholerae* quorum-sensing circuit. (a) Signal transduction at LCD. During this stage, autoinducer levels are low and the CqsS and LuxPQ receptors function as kinases. LuxO is phosphorylated, the Qrr1–4 sRNAs are transcribed, and HapR protein is not produced. (b) Signal transduction at HCD. During this stage, autoinducer levels are high and the CqsS and LuxPQ receptors function as phosphatases. LuxO is unphosphorylated, Qrr1–4 sRNAs are not transcribed, and HapR protein is produced. Solid and dotted lines denote regulatory factors that are produced and not produced, respectively.

genes important for biofilm formation and virulence factor production (32, 146).

Based on sequence homology and circuit configuration, at first analysis it appears that the *V. harveyi* and *V. cholerae* systems are nearly identical, and in turn, should function analogously. However, systematic analyses show striking differences in how the two systems operate. These differences could not have been uncovered through genomic sequence comparisons alone. In the following sections, we compare and contrast these two quorum-sensing systems. First, we outline each system's core components: the multiple autoinducers and their synthases and the molecular mechanisms used by the cognate receptors for signal detection and transduction. Second, we discuss the distinct mechanisms by which the *V. harveyi* and *V. cholerae* Qrr sRNAs function. Third, we discuss how feedback regulation is integrated into the two circuits to uniquely optimize their outputs.

It is interesting to note that many of the quorum-sensing components of the *V. harveyi* and *V. cholerae* systems exist in other *Vibrio* species that harbor LuxIR-type systems. For instance, in addition to the extensively studied LuxIR AHL quorum-sensing system, *V. fischeri* possesses homologs of the *V. harveyi* cascade: LuxMN and LuxS/LuxPQ, as well as LuxU, LuxO, one Qrr sRNA, and a *V. harveyi*-like LuxR called LitR (22). The LuxM and LuxN homologs in *V. fischeri* are called AinS and AinR, respectively (17, 29, 51, 64, 65). The signal produced by AinS is C8HSL (34). Presumably, as with *V. harveyi*, at low autoinducer concentrations, the LuxQ and AinR kinases transfer phosphate through LuxU to LuxO, which activates transcription of the gene encoding the single Qrr sRNA, which prevents production of LitR. Interestingly, LitR activates transcription of *ainS* generating another positive feedback loop in the *V. fischeri* network (64). Moreover, LitR activates transcription of *luxI* linking the LuxS/LuxPQ and AinS/AinR systems to the canonical LuxI/LuxR system (22).

*Vibrio anguillarum*, the causative agent of terminal hemorrhagic septicemia in marine fish, possesses three parallel quorum-sensing

systems homologous to those of *V. harveyi* (14, 15, 78). VanM, the LuxM homolog, produces two autoinducers C6HSL and 3OC6HSL. Both autoinducers are detected by the VanN receptor (15). Unlike *V. harveyi* and *V. cholerae*, the mRNA of *vanT* (the homolog of *luxR/hapR/litR*) is stable at LCD and further production is not induced by quorum sensing (15). It is proposed that inhibition of translation of *vanT* mRNA at LCD occurs through some other unidentified mechanism (77).

### SYNTHESIS OF *V. harveyi* AND *V. cholerae* AUTOINDUCERS

The autoinducers HAI-1, AI-2, and CAI-1 are synthesized by the cytoplasmic enzymes LuxM, LuxS, and CqsA, respectively (3, 37, 38, 75, 105). As mentioned, LuxM has no significant sequence homology to LuxI of *V. fischeri*, but it is similar to another AHL synthase, AinS (29). Mechanistic studies of LuxM synthesis of 3OHC4HSL (**Figure 1a**) have not been performed. However, parallel studies on AinS show that, analogous to the LuxI class of enzymes, the AinS AHL synthase (and presumably the LuxM synthase) also uses SAM and acyl-CoA or acyl-ACP as substrates to produce its particular AHL (34).

The LuxS synthase produces 4,5-dihydroxy-2,3-pentanedione (DPD) (**Figure 1c**), which is the precursor to a set of interconverting molecules that are generically called AI-2. Specifically, DPD is produced from SAM in three enzymatic steps (105). First, methyltransferase enzymes catalyze transfer of the methyl group on SAM to particular substrates to produce products such as DNA, RNA, and proteins. *S*-adenosylhomocysteine (SAH) is formed as a toxic byproduct of these reactions. The Pfs nucleosidase relieves the SAH toxicity by cleaving adenine from SAH to form *S*-ribosylhomocysteine (SRH) (105). SRH is next hydrolyzed by LuxS to form two products: homocysteine and DPD. DPD is unstable and spontaneously converts into different moieties in solution (105). In the marine environment, where the borate concentration

can reach 0.4 mM, DPD cyclizes and reacts with borate to form (2*S*, 4*S*)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran borate (*S*-THMF borate), which is the active form of the AI-2 autoinducer used by *V. harveyi* and *V. cholerae* (**Figure 1c**) (13). In terrestrial environments where boron is limited, (2*R*,4*S*)-2-methyl-2,3,3,4-tetrahydroxytetrahydrofuran (*R*-THMF), an unborated rearranged DPD moiety, is the form of AI-2 used by enteric bacteria such as *Escherichia coli* and *Salmonella typhimurium* (**Figure 1c**) (76). The AI-2 synthase LuxS has gained attention because LuxS homologs exist in hundreds of bacterial genomes, and AI-2 is proposed to be a rather universal signal that fosters interspecies cell-cell communication (21, 142). In *V. harveyi* and *V. cholerae*, and in some other quorum-sensing bacteria, AI-2 clearly acts as a bona fide autoinducer signal (98, 121). Nonetheless, some studies suggest that phenotypes observed in *luxS* mutants of other bacterial species stem from LuxS's role in metabolism of SAM (128). Both possibilities could be correct.

CAI-1 was recently purified from *V. cholerae* and identified as (*S*)-3-hydroxytridecan-4-one (**Figure 1d**) and, as mentioned, its synthesis depends on CqsA, an enzyme with similarity to aminotransferases (38). Purified protein studies show that 3-aminotridecan-4-one (amino-CAI-1), (**Figure 1d**) is in fact the CqsA product and its substrates are (*S*)-2-aminobutyrate and decanoyl-CoA. Crystallographic analysis combined with mutagenesis and spectral analyses verify these findings and demonstrate that *V. cholerae* CqsA produces amino-CAI-1 by a pyridoxal phosphate-dependent aminotransferase reaction. Both amino-CAI-1 and CAI-1 are detected by the *V. cholerae* CqsS receptor, and they have comparable biological activities. However, CAI-1, not amino-CAI-1, is the major form of the molecule in cell-free culture fluids. The current understanding is that, once synthesized, amino-CAI-1 is immediately converted into CAI-1 presumably via another enzyme, and the latter molecule is the predominant form of the signaling molecule encountered by *V. cholerae*. As

---

**SAH:** *S*-adenosylhomocysteine

**SRH:** *S*-ribosylhomocysteine

---

mentioned, *V. harveyi* and several other *Vibrio* species possess homologs of both *cqsA* and *cqsS*, therefore, it is proposed that the CqsA/CqsS system is used for inter-*Vibrio* communication. Consistent with this, cell-free culture fluids from *V. harveyi* and other *Vibrio* species that possess CqsA activate gene expression in a *V. cholerae* CAI-1 reporter strain (37). Whether the CAI-1 autoinducers from *V. harveyi* and other *Vibrio* species are identical to *V. cholerae* CAI-1 or whether they are closely related molecules remains to be addressed. One oddity is that homologs of *cqsA* and *cqsS*, called *lqsA* and *lqsS* respectively, exist in the distantly related bacterium *Legionella pneumophila*. LqsA produces 3-hydroxypentadecan-4-one; a molecule with a longer hydrocarbon chain than CAI-1 (113). The Lqs system promotes host-bacterial interactions in the *L. pneumophila* stationary-phase virulence regulatory network (122, 123).

### SIGNAL TRANSDUCTION THROUGH THE *V. harveyi* AND *V. cholerae* QUORUM-SENSING RECEPTORS

As discussed above, both the *V. harveyi* and *V. cholerae* quorum-sensing systems employ multiple two-component proteins and a phosphorylation-dephosphorylation cascade for signal transduction (37, 75). At LCD, when autoinducer concentration is low, the LuxN, LuxPQ, and CqsS quorum-sensing receptors function as kinases (**Figures 4a** and **5a**). At HCD, when autoinducer concentration is high, the receptors function as phosphatases (**Figures 4b** and **5b**). Although hundreds of two-component systems are known, it is unclear how the phosphorylation activities of particular histidine sensor kinases are regulated by specific signals because only a few ligands have been identified. The *V. harveyi* quorum-sensing system has emerged as an important system for understanding two-component receptor signal transduction across the membrane because the structures of the HAI-1 and AI-2 autoinducers have been known for some time and synthetic molecules are available. These two features made analysis of two-component

signaling across the membrane feasible. Two particular studies, one of AI-2-LuxPQ signaling and one of HAI-1-LuxN signaling, provide insight into how ligand binding elicits the switch in a two-component receptor from kinase to phosphatase (13, 84, 85, 120).

The first informative study concerns AI-2 and LuxPQ. The crystal structures of LuxP in complex with the periplasmic domain of LuxQ (LuxQ<sub>p</sub>) in both the AI-2-free and AI-2-bound forms were solved and compared (84, 85). LuxP, similar to other periplasmic binding proteins, binds AI-2 in a cleft formed between two similar domains connected by a three-stranded hinge. LuxQ<sub>p</sub> is composed of two tandem PAS domains with no sequence homology to one another or to other PAS folds. Unliganded LuxP adopts an open conformation representing the AI-2 receptive state in the apo-LuxP/LuxQ<sub>p</sub> complex. Mutations that destabilize or eliminate the interfaces between LuxP and LuxQ<sub>p</sub> decrease the concentration of AI-2 required to convert LuxQ from kinase to phosphatase, suggesting that interactions between unliganded LuxP and LuxQ<sub>p</sub> inhibit the conversion of LuxQ from kinase to phosphatase. Binding of AI-2 to LuxPQ<sub>p</sub> induces significant conformational and organizational changes in the complex. First, the two domains in the AI-2-bound LuxP close via a dramatic conformational change. This event, however only minimally alters the LuxQ<sub>p</sub> conformation. What is critical is that, AI-2 binding alters LuxPQ<sub>p</sub>-LuxPQ<sub>p</sub> dimerization with the two LuxPQ<sub>p</sub> dimers undergoing an approximately 140 degree rotation around an axis between them. Mutations constructed to investigate the role of the new interface formed between the LuxPQ<sub>p</sub>-LuxPQ<sub>p</sub> dimers when AI-2 binds showed that they decrease responsiveness to AI-2, suggesting that the mutations inhibit the two LuxPQ<sub>p</sub> dimers from making the interface required to promote phosphatase activity. Therefore, increased AI-2 is required to switch LuxQ to phosphatase mode. The current model proposes that two LuxPQ complexes form a symmetric heterotetramer (LuxPQ-LuxPQ) in the absence of AI-2 at LCD. Analogous to what has been

shown for the histidine kinases EnvZ and NtrB (86, 143), autophosphorylation is presumed to occur by cytoplasmic cross-phosphorylation between two LuxQ histidine kinase monomers (although this has not been verified for LuxQ). At HCD, AI-2 binding to LuxP causes a large rotation of one of the LuxPQ subunits relative to the other LuxPQ subunit. This rotational change disrupts the symmetry of the LuxPQ-LuxPQ tetramer, and thus prevents cross-phosphorylation between the cytoplasmic regions of the two LuxQ monomers. Moreover, the crystal structures predict that this asymmetric architecture prevents formation of higher order oligomers, suggesting that AI-2-bound LuxPQ tetramers cannot cluster (85). This arrangement potentially reduces premature commitment to quorum sensing as a consequence of signal noise (see below).

The second informative signaling study concerns LuxN. The proposed autoinducer binding domain of LuxN contains nine transmembrane-spanning (TM) helices with the N-terminus located on the periplasmic side of the inner membrane (47). Because of this topology, no detailed structural information is available. Rather, an approach that exploited mutagenesis and suppressor analyses, together with HAI-1 antagonist studies and mathematical modeling, was used to define the HAI-1 binding site and the *in vivo* signaling parameters of the LuxN receptor (120). Mutations in TMs 4, 5, 6, and 7, and the intervening periplasmic loops 2 and 3 of LuxN, render *V. harvey* nonresponsive to HAI-1. Analyses of the mutants showed that one class does not bind HAI-1 and thus does not switch from kinase to phosphatase. A second class of mutants has reduced affinity for HAI-1 and thus can switch from kinase to phosphatase albeit only at high concentrations of HAI-1.

A potent competitive HAI-1 antagonist was identified from a high-throughput chemical screen (120). This antagonist was used to further probe the LuxN/HAI-1 interaction. By simultaneously varying the amounts of antagonist and HAI-1, HAI-1 dose-response curves could be generated, which first, defined

the HAI-1  $EC_{50}$  value for wild-type LuxN to be 20 nM. Second, the data from all the curves could be collapsed into a single curve. The principle underlying the data fitting is that there is a fixed relation between the kinase and phosphatase configurations of LuxN such that the probability for LuxN to be a kinase depends on the free energy difference between the two configurations. Using this analysis, the dissociation constants ( $K_D$ ) for HAI-1 for wild-type LuxN and the various LuxN mutants were determined. In the phosphatase state,  $K^{HAI-1}$  is approximately 1 nM, and in the kinase state,  $K^{antagonist}$  is approximately 500 nM. The mutants could then be classified as those that affect HAI-1 binding (i.e., with altered  $K_D$ ), and those that do not affect HAI-1 binding (similar  $K_D$ ) but have altered free energy differences between the two configurations (kinase and phosphatase) of LuxN. One additional prediction from these analyses is that the probability for LuxN to be a kinase in the absence of HAI-1 is approximately 96%, which explains the large ratio between the  $EC_{50}$  value (20 nM) and the underlying  $K_D$  (1 nM).

This final observation of a large difference in ligand  $EC_{50}$  and  $K_D$  differs dramatically from what has been found in the classic chemotaxis two-component signaling network where there exists only a small difference ( $\sim 0.1$ ) between  $EC_{50}$  and  $K_D$ . Thus, in the chemotaxis system, there is a roughly equal probability for the receptor to be a kinase or a phosphatase when ligand concentration is low (50). The difference between the receptors in the quorum-sensing and chemotaxis circuits apparently allows each system to solve its respective biological problem effectively. In chemotaxis, bacteria need to respond rapidly to small alterations in signal concentrations, therefore, the receptors are poised to immediately change from kinase to phosphatase and vice versa by spending half of the time in each state. Furthermore, chemotaxis receptors are clustered in arrays that promote amplification of the signal (68, 112, 115). By contrast, quorum-sensing receptors, as discussed above, do not cluster and distribute evenly on the bacterial inner membrane (85),

so they do not amplify small perturbations in ligand concentration. Furthermore, switching from kinase to phosphatase requires a significant buildup of autoinducer given that the receptors spend almost 100% of the time in the kinase state (120). An overall reduced signal sensitivity in the quorum-sensing system presumably prevents accidental commitment to group behavior in response to signal noise.

Although each of the above studies was focused exclusively on one *V. harveyi* receptor—structural analysis of LuxPQ and antagonism analysis of LuxN—it is assumed that the intrinsic signaling properties (i.e., the nonclustering of receptors observed for LuxPQ and the low signal sensitivity observed for LuxN) will apply generally to the other *V. harveyi* and *V. cholerae* quorum-sensing receptors (e.g., *V. harveyi* and *V. cholerae* CqsS, *V. cholerae* LuxPQ). In fact, studies on *V. cholerae* suggest that the CqsS receptor also has low signal sensitivity (W.-L. Ng, unpublished observation).

### **FUNCTIONS OF Qrr sRNAs IN *V. harveyi* AND *V. cholerae*: ADDITIVITY VERSUS REDUNDANCY**

At the heart of *V. harveyi* and *V. cholerae* quorum-sensing circuits lies multiple Qrr sRNAs, and the precisely controlled levels of these sRNAs dictate whether cells switch into or out of quorum-sensing behavior (Figures 4 and 5). At LCD, the Qrr sRNAs are transcribed and prevent production of LuxR/HapR. Qrr sRNAs function by Hfq-assisted base pairing with the mRNA of *luxR/hapR*, which blocks the translation and destabilizes the transcript (57, 124). At HCD, the Qrr sRNAs are not transcribed, the *luxR/hapR* mRNA accumulates and LuxR/HapR protein is produced. As mentioned above, *V. harveyi* possess five Qrr sRNAs (Qrr1–5) and *V. cholerae* possess four Qrr sRNAs (Qrr1–4). The LCD transcription of the Qrr sRNAs is controlled by LuxO-P in both *V. harveyi* and *V. cholerae*. In *V. harveyi*, the steady state level at LCD is highest for Qrr4, followed by Qrr2, Qrr3, Qrr1, and then Qrr5 (124). In *V. cholerae*, a similar pattern is observed (i.e.,

Qrr4 > Qrr2 ≈ Qrr3 > Qrr1) (56, 57, 118). Because of their unique expression levels, it is presumed that the relative strength of each sRNA in controlling quorum-sensing regulated genes via LuxR/HapR follows the same order. Although the sequences of the Qrr sRNAs required for targeting the *luxR/hapR* mRNAs are identical in both *V. harveyi* and *V. cholerae* and the overall sequences of the mRNAs are extraordinarily similar (>80% identity), the sRNAs nonetheless function by different means in the two *Vibrio* species.

In *V. cholerae*, Qrr1–4 function redundantly to regulate quorum sensing. That is, any single Qrr sRNA is sufficient to destabilize the *hapR* mRNA and prevent production of HapR. Therefore, the simultaneous deletion of the four *qrr* sRNA genes is required to alter quorum sensing (57). Functional redundancy of the *V. cholerae* four Qrr sRNAs stems from *qrr* gene dosage compensation. Transcription of a particular *qrr* gene is affected by the amount of the other three Qrr sRNAs present (118). For example, in the absence of three Qrr sRNAs, transcription of the remaining *qrr* sRNA gene is increased. *qrr* gene dosage compensation maintains the Qrr sRNA pool within a specific range. Dosage compensation depends on two feedback loops called the HapR-Qrr feedback loop and the LuxO-Qrr feedback loop, which are discussed below.

In stark contrast to how the Qrrs function in *V. cholerae*, in *V. harveyi* the five Qrr sRNAs work additively to control quorum sensing. Thus, deletion of any single *qrr* gene results in a quorum-sensing phenotypic change (124). As mentioned, the strength of each Qrr sRNA in controlling quorum sensing mirrors its respective expression level (i.e., Qrr4 > Qrr2 > Qrr3 > Qrr1 > Qrr5). Therefore, quorum-sensing behavior is nearly wild type in a mutant possessing only *qrr4*, whereas in a mutant possessing only *qrr5*, quorum-sensing behavior is nearly nonexistent (124). The two feedback loops analogous to those in *V. cholerae*, in this case called the LuxR-Qrr feedback loop and the LuxO-Qrr feedback loop, also exist and function in *V. harveyi* (124a, 125). For

reasons that are not yet clear, *V. harveyi* Qrr sRNA mutants do not fully dosage compensate, which results in the observed alterations in their quorum-sensing phenotypes. One hypothesis is that transcription of each *V. harveyi* *qrr* gene is subject to additional controls that allow fine-tuning of LuxR production under different environmental conditions in addition to the autoinducer inputs.

## PROCESSING THE INFORMATION CONTAINED IN MULTIPLE AUTOINDUCERS IN *V. harveyi* AND *V. cholerae*

It is not uncommon for a single bacterial species to produce and detect multiple autoinducers (AHLs or peptides) and/or to produce more than one type of autoinducer (44, 91, 110, 111, 139). In these cases, unique information is contained in each autoinducer, and the bacterium has some mechanism to differentiate between and discretely respond to each signal. For example, *P. aeruginosa* produces two AHLs, 3OC12-HSL and C4-HSL, by LasI and RhlI, respectively (**Figure 1a**). The systems are arranged in series such that expression of the Rhl system is activated by the LasR-AHL bound complex. Hence, production of 3OC12HSL precedes production of C4HSL (52, 72, 93); and likewise 3OC12HSL-responsive genes are expressed prior to those responsive to C4HSL. *P. aeruginosa* also produces a third autoinducer, 2-heptyl-3-hydroxy-4-quinolone, designated the *Pseudomonas* quinolone signal (PQS) (**Figure 1e**) (92). Production of PQS is influenced both positively and negatively by the LasIR and RhlIR systems, respectively (71, 132). In *B. subtilis*, the peptide autoinducers ComX and CSF have opposing functions in the control of the competence and sporulation pathways (69, 111). ComX accumulation stimulates ComP-dependent phosphorylation of the ComA response regulator, which promotes competence (69). By contrast, a high concentration of CSF antagonizes ComX-induced phosphorylation of ComA, which decreases competence development, and instead favors

the sporulation pathway (53). As mentioned, four *S. aureus* specificity groups exist, and each group produces an autoinducer peptide that functions as a quorum-sensing agonist in its own *S. aureus* group but acts as a quorum-sensing antagonist in the heterologous *S. aureus* groups, resulting in interference in the latter's quorum-sensing response (42, 44, 66).


Surprisingly, the sensory information contained in the three *V. harveyi* autoinducers, and likewise in the two *V. cholerae* autoinducers, is transduced into the cells via shared phosphorelay cascades. Specifically, in both *Vibrio* species, the two-component autoinducer receptors channel phosphate to and from a single phosphotransfer protein LuxU (see **Figures 4 and 5**). This network arrangement raises the intriguing question of whether or not *V. harveyi* and *V. cholerae* can distinguish between the different autoinducers. Although this is not completely understood at present, it is clear that, at a minimum, the different autoinducer signals certainly have different strengths. In *V. harveyi*, the HAI-1 signal is stronger than AI-2, which is stronger than CAI-1 (37). By contrast, in *V. cholerae*, CAI-1 is stronger than AI-2 (75). The differences in signal strengths are likely due to differences in the relative enzymatic activities of the respective receptor kinases/phosphatases. Differences in the intrinsic receptor signaling parameters (e.g., binding constants, free energy differences between on/off configurations) may also play roles in the strengths of the different autoinducers.

The question of the mechanism underlying signal discrimination in the *Vibrios* is being most intensively investigated in *V. harveyi* with respect to how the bacterium distinguishes between its two strongest signals, HAI-1 and AI-2 (60, 80, 134). Several studies, although not yet conclusively solving the problem, have provided insight into how a bacterium can use shared regulatory components and nonetheless respond discretely to the different signals. In one study, expression of more than 50 *V. harveyi* quorum-sensing regulated genes was analyzed in the presence of only HAI-1, only AI-2, or both autoinducers together. Three

**GFP:** green fluorescent protein

classes of genes were identified: Class 1 genes show significant regulation only in the simultaneous presence of HAI-1 and AI-2 (i.e., coincidence regulation); Class 2 genes exhibit an alteration in expression when either HAI-1 or AI-2 is present, and expression changes more drastically when both autoinducers are supplied together; Class 3 genes exhibit expression changes in the presence of either HAI-1 alone, or AI-2 alone, and supplying both autoinducers simultaneously does not produce a response that is different from one autoinducer alone (134). This report demonstrated that the various combinations of autoinducers produce graded but inversely correlated changes in *Qrr* sRNA and *LuxR* concentrations (Figure 6). Specifically, low concentrations of autoinducers (i.e., at LCD) lead to high *qrr* and low *luxR* expression. High concentrations of autoinducers (i.e., at HCD) result in low *qrr* and high *luxR* expression (See Reference 134 and Figures 4 and 6). Thus, the three classes of target-gene responses can be understood in terms of a *LuxR* promoter affinity model (134). Target genes with promoters that have high affinity for *LuxR* are regulated in response to the lowest concentrations of autoinducer, and thus these genes (Class 3) are the first ones activated/repressed when cells switch into quorum-sensing mode. Genes with promoters that have low affinity for *LuxR* respond to only the highest autoinducer concentrations and are regulated at later times in growth (Class 1). Class 2 genes, which respond to mid-level autoinducer concentrations, are expressed after Class 3 and prior to Class 1 genes (See Reference 134 and Figure 6).

To further understand how *LuxR* regulates genes with different affinities, the *LuxR* DNA recognition sequence was identified using protein binding microarrays (97). The consensus sequence for *LuxR* binding contains a 21 bp operator with dyad symmetry, and the critical bases for binding in each half-site are independent of one another. Some *LuxR*-regulated genes possess multiple binding sites in their promoters, suggesting cooperative binding which may also play a role in the timing and strength of *LuxR*-dependent regulation



**Figure 6**

Reciprocal production of *V. harveyi* *Qrr* sRNAs and *LuxR* leads to temporal control of quorum-sensing target genes. From LCD to HCD, *Qrr* sRNA concentrations decrease and *LuxR* concentrations increase. As a consequence, Class 3 quorum-sensing target genes, whose promoters have the highest affinity for *LuxR*, are activated/repressed first, followed by Class 2 genes, and finally Class 1 genes.

(97). Another study with SmcR, the *LuxR* homolog in *Vibrio vulnificus*, defined a 22 bp consensus sequence that is highly similar to that found for *V. harveyi* *LuxR* (54).

Using single-cell fluorescence microscopy analyses of *qrr-gfp*, the *V. harveyi* responses to HAI-1 and AI-2 were quantified. Each autoinducer contributes nearly equally to the total output response (60). Thus, the information from the two distinct autoinducers is combined additively. Based on these analyses, it is proposed that *V. harveyi* can distinguish between at least three distinct conditions of external autoinducer. First, high *qrr-gfp* expression occurs when both HAI-1 and AI-2 concentrations are low. Second, low *qrr-gfp* expression occurs when both HAI-1 and AI-2 concentrations are high. Third, intermediate *qrr-gfp* expression occurs when one autoinducer concentration is low and the other is high. However, in this final situation, high HAI-1 combined with low AI-2 concentration is indistinguishable from low HAI-1 combined with high AI-2 concentration (60). These recent findings led to the idea that detecting multiple autoinducers allows *V. harveyi* to monitor the developmental stage of the community. This model assumes that production of each autoinducer follows an invariant temporal order in which high HAI-1/low AI-2 and low HAI-1/high AI-2 are mutually



exclusive (60). These findings also suggest that autoinducers allow populations of *V. harveyi* to monitor the species composition of the community. As mentioned, the different autoinducers specify the relatedness of the members of the community: HAI-1 is only produced by *V. harveyi*, CAI-1 is produced by many *Vibrios*, and AI-2 is produced by widely diverse bacterial species. Therefore, the different combinations of autoinducers could reflect the composition and abundance of species in the vicinal community (60). Because the *V. harveyi* CqsA/CqsS system has not yet been included in these types of investigations and no signal discrimination studies have been performed in *V. cholerae*, it remains an open question as to whether these findings can be more generally applied.

### FEEDBACK CONTROL OF QUORUM SENSING IN *V. harveyi* AND *V. cholerae*

Positive feedback is a hallmark of quorum-sensing regulatory networks. As mentioned, in canonical LuxI/LuxR quorum-sensing systems, expression of the autoinducer synthase gene is positively controlled by the AHL-bound LuxR type protein (18, 26, 108). The outcome of these feedback loops is accelerated production of AHL autoinducer, which leads to synchrony in quorum-sensing behavior. Positive feedback loops exist in oligopeptide-based Gram-positive quorum-sensing systems (45, 95). In many cases, the genes encoding the peptide signal, the histidine kinase receptor, the cognate response regulator, and its accessory factors form an operon (for example, the *agrBDCA* and *comCDE* operons of *S. aureus* and *S. pneumoniae*, respectively). Furthermore, typically the response regulator in the system acts as an auto-activator of the operon. This autoregulatory wiring fosters positive feedback through which the amounts of the peptide ligand, the membrane receptor, and the response regulator all increase drastically once the autoinducer has accumulated above the initial concentration required for detection.

In *V. harveyi* and *V. cholerae*, a different set of feedback loops have recently been identified

that ensure precise timing of quorum-sensing transitions (Figure 7). These feedback loops are summarized as follows:

1. HapR/LuxR auto-repression loop
2. HapR/LuxR-Qrr feedback loop
3. LuxO auto-repression loop
4. LuxO-Qrr feedback loop.

#### HapR/LuxR Auto-Repression Loop


HapR and LuxR bind to their own promoters and repress transcription (12, 59), which results in a steady increase in HapR/LuxR concentrations as cell density increases. Thus, this feedback loop, by preventing runaway expression of LuxR/HapR, minimizes the chances of premature commitment of the cells to population-wide changes in gene expression.

#### HapR/LuxR-Qrr Feedback Loop

During the transition from LCD to HCD, LuxO-P concentrations decrease as autoinducer concentrations increase, resulting in decreased *qrr* transcription and, in turn, increased HapR/LuxR production (Figures 4 and 5). HapR/LuxR feeds back as a transcriptional activator of the *qrr* genes (Figure 7) (119, 125). When cells switch from LCD to HCD, the HapR/LuxR-Qrr feedback loop prolongs the production of Qrr sRNAs and delays the entry into HCD mode. By contrast, when cells switch from HCD to LCD, the HapR/LuxR-Qrr feedback loop dramatically increases expression of the *qrr* genes and accelerates the transition out of quorum-sensing mode and into individual-cell behavior (119, 125). Interestingly, HapR (*V. cholerae*) acts indirectly on *qrr1*–4 in the HapR-Qrr feedback loop, whereas LuxR (*V. harveyi*) binds directly to *qrr* promoters in the feedback mechanism. However, in *V. harveyi*, *qrr2*, *qrr3*, and *qrr4* are subject to LuxR feedback control but *qrr1* and *qrr5* are not (119, 125).

#### LuxO Auto-Repression Loop

The *qrr1* gene and the *luxO* gene lie adjacent to one another in the genome and are transcribed



**Figure 7**

Feedback loops identified in the *V. harveyi* and *V. cholerae* quorum-sensing networks. Four different feedback loops are integrated into the *V. harveyi* and *V. cholerae* quorum-sensing circuits. Arrows denote activation. T-shape arrows denote repression.

divergently (57). The LuxO-binding site required for *qrr1* expression overlaps with the  $-35$  site in the *luxO* promoter. This unique organization allows LuxO to simultaneously activate expression of *qrr1* and repress its own transcription via blocking access to RNAP for transcription (124a). Although LuxO requires phosphorylation to act as a transcriptional activator, LuxO-auto-repression does not require that LuxO be phosphorylated (118). The LuxO auto-repression loop limits the amount of LuxO to within a narrow window, the consequence of which is to carefully control production of the Qrr sRNAs. The LuxO auto-repression loop is critical for ensuring precise timing of the quorum-sensing transition (124a).

### LuxO-Qrr Feedback Loop

Analogous to the way the Qrr sRNAs act on the *luxR/hapR* mRNAs, the Qrr sRNAs bind to and destabilize the mRNA encoding LuxO and stimulate its degradation. This prevents LuxO protein production. The LuxO-Qrr feedback loop works synergistically with the LuxO auto-repression loop, to restrict Qrr sRNA levels to only a narrow range by limiting fluctuations in LuxO levels (118, 124a). Also, as discussed above, the other function of the LuxO-Qrr feedback loop is in *qrr* gene dosage compensation (118).

Why are so many feedback loops involved in controlling the *V. harveyi* and *V. cholerae* quorum-sensing responses? As discussed above, the quorum-sensing receptors are insensitive to small perturbations in signal suggesting that the *V. harveyi* and *V. cholerae* networks are tuned to ignore sudden fluctuations in the environment. Presumably then, a sudden surge in any component in the quorum-sensing network would be detrimental. Indeed, all of the studies of the *V. harveyi* and *V. cholerae* feedback loops point to their functioning together to minimize fluctuations in the levels of individual components in the quorum-sensing circuits. The LuxR/HapR auto-repression loop prevents sudden increases in the level of the master quorum-sensing regulator LuxR/HapR (12, 59). The LuxR/HapR-Qrr feedback loop also delays production of LuxR/HapR (119, 125). Finally, the LuxO-Qrr feedback loop together with the LuxO auto-repression loop limit fluctuations in LuxO levels, which prevents surges of Qrr sRNA levels (118, 124a). We note that feedback loops similar to those described in these two quorum-sensing systems are common network motifs in biological circuits (99). Feedback loops are known to reduce variations in the steady-state levels of regulatory components. In the case of the *V. harveyi* and *V. cholerae* quorum-sensing networks, by minimizing alterations in cytoplasmic quorum-sensing

**RNAP:** RNA polymerase

regulatory components, the feedback loops, in turn, limit cell-to-cell variations in behavior. In so doing, the feedback loops impose synchrony on the population-wide quorum-sensing response which is imperative for successful collective behaviors.

## OUTLOOK AND FUTURE DIRECTIONS

A major goal of studying the quorum-sensing systems of *V. harveyi* and *V. cholerae* is to understand at the molecular, cellular, and population levels the process used by bacteria for cell-cell communication. Fundamental questions pertaining to this goal remain. Multiple quorum-sensing signals are channeled into a single circuit; yet different signal inputs lead to differential gene expression outputs. Particularly critical is to understand what governs the relative signal strengths of LuxM/LuxN, LuxS/LuxPQ, and CqsA/CqsS. How does CqsS detect and respond to both CAI-1 and amino-CAI-1, and do LuxPQ and CqsS receptors have signaling parameters similar to those of LuxN? Moreover, the networks employ both RNA-based and protein-based regulatory factors. Until recently, sRNA-mediated gene regulation was underappreciated in bacteria (30, 135). Thus, it remains to be investigated why sRNAs are optimal for quorum-sensing regulation and what advantages multiple sRNAs provide the circuit. For instance, are sRNA regulators more precise than protein regulators in controlling gene expression because they are less prone to fluctuation? Do Qrr sRNAs have different affinities for their targets [e.g., *luxO*, *luxR/bapR*, and the recently discovered *vca0939* (33)], and if so, does this impinge on the dynamics of the quorum-sensing transitions? What

is the molecular mechanism underpinning the differential expression of the *qrr* genes? What is probably most remarkable when one ponders the *V. harveyi* and *V. cholerae* quorum-sensing systems is that they employ a set of nearly identical constituent components, yet the functioning of these components and the behaviors of the two systems are dramatically different.

Finally, many open questions remain regarding the evolutionary forces that shape quorum-sensing systems and how quorum-sensing behaviors of individual cells translate into the collective properties of bacterial groups (82, 137). Recent work has shown that populations of *P. aeruginosa*, which use quorum sensing to up-regulate virulence at HCD, can be invaded both by mutants that do not produce autoinducers and by mutants that do not respond to autoinducers (16, 102). When mixed with wild-type cells in equal proportion, exploitative *P. aeruginosa* quorum-sensing mutants decreased the virulence profiles of both acute and chronic *P. aeruginosa* infections in a mouse model (101). In contrast to *Pseudomonas* and many other systems, the *Vibrios* activate virulence factor expression at LCD and repress these traits at HCD. Theory suggests that *Vibrios* do so in order to efficiently escape from their hosts (83, 146), but these predictions have not yet been tested by experiments. These newest findings make it clear that the evolution of quorum sensing on short time scales must be addressed as the field turns to the development of biotechnological therapies to manipulate quorum sensing. Furthermore, clarifying the ecological pressures that favor particular quorum-sensing regulatory strategies may allow us to better understand how quorum sensing has evolved and continues to evolve in bacterial populations.

### HIGHLIGHTS

1. Bacteria use extracellular chemical signal molecules called autoinducers for quorum sensing. Quorum sensing is a cell-cell communication process used to monitor cell number and species complexity in a population. The *V. harveyi* and *V. cholerae* quorum-sensing networks exhibit similarity to both canonical Gram-negative and Gram-positive quorum-sensing systems.

2. *V. harveyi* possesses three quorum-sensing systems (LuxM/LuxN, LuxS/LuxPQ, and CqsA/CqsS). *V. cholerae* possesses two quorum-sensing systems (LuxS/LuxPQ, CqsA/CqsS). Although components of the two systems are similar, distinctive features greatly alter their biology.
3. The quorum-sensing receptors in *V. harveyi* and *V. cholerae* are organized such that they prevent dramatic changes in response to small perturbations in autoinducer signals. This arrangement stands in contrast to the bacterial chemotaxis receptor system, which is exquisitely sensitive to small changes in ligand concentration.
4. Small RNAs (sRNAs) lie at the core of both the *V. harveyi* and *V. cholerae* quorum-sensing systems. The sRNAs function differently in these two species. In *V. harveyi*, five sRNAs function additively. In *V. cholerae*, four sRNAs function redundantly.
5. Multiple autoinducers are detected and integrated through shared phosphorelay systems in both the *V. harveyi* and *V. cholerae* quorum-sensing systems. Differences exist in the strength of each signal.
6. Multiple feedback loops exist in the *V. harveyi* and *V. cholerae* quorum-sensing networks. These feedback loops prevent fluctuations in the amounts of regulatory components in the respective systems, and ensure precise input-output relationships.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

We thank members of the Bassler laboratory for insightful discussions and Mr. Carey Nadell for his input on evolution of group behaviors in bacteria. This work was supported by NRSA postdoctoral fellowship F32GM082061 for W-L Ng; HHMI, NIH grants RO1GM065859 and RO1AI054442, and NSF grant MCB0639855 for B. L. Bassler.

## LITERATURE CITED

1. Albus AM, Pesci EC, Runyen-Janecky LJ, West SE, Iglewski BH. 1997. Vfr controls quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* 179:3928–35
2. Austin B, Zhang XH. 2006. *Vibrio harveyi*: a significant pathogen of marine vertebrates and invertebrates. *Lett. Appl. Microbiol.* 43:119–24
3. Bassler BL, Wright M, Showalter RE, Silverman MR. 1993. Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence. *Mol. Microbiol.* 9:773–86
4. Bassler BL, Wright M, Silverman MR. 1994. Multiple signalling systems controlling expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a second sensory pathway. *Mol. Microbiol.* 13:273–86
5. Bassler BL, Wright M, Silverman MR. 1994. Sequence and function of LuxO, a negative regulator of luminescence in *Vibrio harveyi*. *Mol. Microbiol.* 12:403–12
6. Blokesch M, Schoolnik GK. 2007. Serogroup conversion of *Vibrio cholerae* in aquatic reservoirs. *PLoS Pathog* 3:e81
7. Blokesch M, Schoolnik GK. 2008. The extracellular nuclease Dns and its role in natural transformation of *Vibrio cholerae*. *J. Bacteriol.* 190:7232–40

8. Bottomley MJ, Muraglia E, Bazzo R, Carfi A. 2007. Molecular insights into quorum sensing in the human pathogen *Pseudomonas aeruginosa* from the structure of the virulence regulator LasR bound to its autoinducer. *J Biol Chem* 282:13592–600
9. Callahan SM, Dunlap PV. 2000. LuxR- and acyl-homoserine-lactone-controlled non-lux genes define a quorum-sensing regulon in *Vibrio fischeri*. *J. Bacteriol.* 182:2811–22
10. Cao JG, Meighen EA. 1989. Purification and structural identification of an autoinducer for the luminescence system of *Vibrio harveyi*. *J. Biol. Chem.* 264:21670–76
11. Case RJ, Labbate M, Kjelleberg S. 2008. AHL-driven quorum-sensing circuits: their frequency and function among the Proteobacteria. *Isme. J.* 2:345–49
12. Chatterjee J, Miyamoto CM, Meighen EA. 1996. Autoregulation of *luxR*: the *Vibrio harveyi lux*-operon activator functions as a repressor. *Mol. Microbiol.* 20:415–25
13. Chen X, Schauder S, Potier N, Van Dorsselaer A, Pelczar I, et al. 2002. Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* 415:545–49
14. Croxatto A, Chalker VJ, Lauritz J, Jass J, Hardman A, et al. 2002. VanT, a homologue of *Vibrio harveyi* LuxR, regulates serine, metalloprotease, pigment, and biofilm production in *Vibrio anguillarum*. *J. Bacteriol.* 184:1617–29
15. Croxatto A, Pride J, Hardman A, Williams P, Camara M, Milton DL. 2004. A distinctive dual-channel quorum-sensing system operates in *Vibrio anguillarum*. *Mol. Microbiol.* 52:1677–89
16. Diggle SP, Griffin AS, Campbell GS, West SA. 2007. Cooperation and conflict in quorum-sensing bacterial populations. *Nature* 450:411–14
17. Dunlap PV. 1999. Quorum regulation of luminescence in *Vibrio fischeri*. *J. Mol. Microbiol. Biotechnol.* 1:5–12
18. Engebrecht J, Nealson K, Silverman M. 1983. Bacterial bioluminescence: isolation and genetic analysis of functions from *Vibrio fischeri*. *Cell* 32:773–81
19. Engebrecht J, Silverman M. 1984. Identification of genes and gene products necessary for bacterial bioluminescence. *Proc. Natl. Acad. Sci. USA* 81:4154–58
20. Faruque SM, Albert MJ, Mekalanos JJ. 1998. Epidemiology, genetics, and ecology of toxigenic *Vibrio cholerae*. *Microbiol. Mol. Biol. Rev.* 62:1301–14
21. Federle MJ, Bassler BL. 2003. Interspecies communication in bacteria. *J. Clin. Invest.* 112:1291–99
22. Fidopiastis PM, Miyamoto CM, Jobling MG, Meighen EA, Ruby EG. 2002. LitR, a new transcriptional activator in *Vibrio fischeri*, regulates luminescence and symbiotic light organ colonization. *Mol. Microbiol.* 45:131–43
23. Freeman JA, Bassler BL. 1999. A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. *Mol. Microbiol.* 31:665–77
24. Freeman JA, Lilley BN, Bassler BL. 2000. A genetic analysis of the functions of LuxN: a two-component hybrid sensor kinase that regulates quorum sensing in *Vibrio harveyi*. *Mol. Microbiol.* 35:139–49
25. Fuqua C, Parsek MR, Greenberg EP. 2001. Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu. Rev. Genet.* 35:439–68
26. Fuqua WC, Winans SC. 1994. A LuxR-LuxI type regulatory system activates *Agrobacterium* Ti plasmid conjugal transfer in the presence of a plant tumor metabolite. *J. Bacteriol.* 176:2796–806
27. Geisinger E, George EA, Muir TW, Novick RP. 2008. Identification of ligand specificity determinants in AgrC, the *Staphylococcus aureus* quorum-sensing receptor. *J. Biol. Chem.* 283:8930–38
28. Geisinger E, Muir TW, Novick RP. 2009. *agr* receptor mutants reveal distinct modes of inhibition by staphylococcal autoinducing peptides. *Proc. Natl. Acad. Sci. USA* 106:1216–21
29. Gilson L, Kuo A, Dunlap PV. 1995. AinS and a new family of autoinducer synthesis proteins. *J. Bacteriol.* 177:6946–51
30. Gottesman S. 2004. The small RNA regulators of *Escherichia coli*: roles and mechanisms. *Annu. Rev. Microbiol.* 58:303–28
31. Gould TA, Schweizer HP, Churchill ME. 2004. Structure of the *Pseudomonas aeruginosa* acyl-homoserinylactone synthase LasI. *Mol. Microbiol.* 53:1135–46
32. Hammer BK, Bassler BL. 2003. Quorum sensing controls biofilm formation in *Vibrio cholerae*. *Mol. Microbiol.* 50:101–4

---

38. The CAI-1 autoinducer from *V. cholerae* was purified and its chemical structure was determined in this study.

---

33. Hammer BK, Bassler BL. 2007. Regulatory small RNAs circumvent the conventional quorum sensing pathway in pandemic *Vibrio cholerae*. *Proc. Natl. Acad. Sci. USA* 104:11145–49
34. Hanzelka BL, Parsek MR, Val DL, Dunlap PV, Cronan JE, Jr, Greenberg EP. 1999. Acylhomoserine lactone synthase activity of the *Vibrio fischeri* AinS protein. *J. Bacteriol.* 181:5766–70
35. Havarstein LS, Coomaraswamy G, Morrison DA. 1995. An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. *Proc. Natl. Acad. Sci. USA* 92:11140–44
36. Henke JM, Bassler BL. 2004. Quorum sensing regulates type III secretion in *Vibrio harveyi* and *Vibrio parahaemolyticus*. *J. Bacteriol.* 186:3794–805
37. Henke JM, Bassler BL. 2004. Three parallel quorum-sensing systems regulate gene expression in *Vibrio harveyi*. *J. Bacteriol.* 186:6902–14
38. Higgins DA, Pomianek ME, Kraml CM, Taylor RK, Semmelhack MF, Bassler BL. 2007. The major *Vibrio cholerae* autoinducer and its role in virulence factor production. *Nature* 450:883–86
39. Hoch JA, Silhavy TJ. 1995. *Two-Component Signal Transduction*. Washington, DC: ASM Press. 488 pp.
40. Hwang I, Li PL, Zhang L, Piper KR, Cook DM, et al. 1994. TraI, a LuxI homologue, is responsible for production of conjugation factor, the Ti plasmid *N*-acylhomoserine lactone autoinducer. *Proc. Natl. Acad. Sci. USA* 91:4639–43
41. Inouye M, Dutta R. 2003. *Histidine kinases in signal transduction*. London: Academic. 520 pp.
42. Jarraud S, Lyon GJ, Figueiredo AM, Gerard L, Vandenesch F, et al. 2000. Exfoliatin-producing strains define a fourth *agr* specificity group in *Staphylococcus aureus*. *J. Bacteriol.* 182:6517–22
43. Jensen RO, Winzer K, Clarke SR, Chan WC, Williams P. 2008. Differential recognition of *Staphylococcus aureus* quorum-sensing signals depends on both extracellular loops 1 and 2 of the transmembrane sensor AgrC. *J. Mol. Biol.* 381:300–9
44. Ji G, Beavis R, Novick RP. 1997. Bacterial interference caused by autoinducing peptide variants. *Science* 276:2027–30
45. Ji G, Beavis RC, Novick RP. 1995. Cell density control of staphylococcal virulence mediated by an octapeptide pheromone. *Proc. Natl. Acad. Sci. USA* 92:12055–59
46. Jobling MG, Holmes RK. 1997. Characterization of *hapR*, a positive regulator of the *Vibrio cholerae* HA/protease gene *hap*, and its identification as a functional homologue of the *Vibrio harveyi luxR* gene. *Mol. Microbiol.* 26:1023–34
47. Jung K, Odenbach T, Timmen M. 2007. The quorum-sensing hybrid histidine kinase LuxN of *Vibrio harveyi* contains a periplasmically located N terminus. *J. Bacteriol.* 189:2945–48
48. Kaplan HB, Greenberg EP. 1985. Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *J. Bacteriol.* 163:1210–14
49. Kelly RC, Bolitho ME, Higgins DA, Lu W, Ng W-L et al. 2009. The *Vibrio cholerae* quorum-sensing autoinducer CAI-1: analysis of the biosynthetic enzyme CqsA. *Nature Chem. Biol.* In press
50. Keymer JE, Endres RG, Skoge M, Meir Y, Wingreen NS. 2006. Chemosensing in *Escherichia coli*: two regimes of two-state receptors. *Proc. Natl. Acad. Sci. USA* 103:1786–91
51. Kuo A, Callahan SM, Dunlap PV. 1996. Modulation of luminescence operon expression by *N*-octanoyl-L-homoserine lactone in *ainS* mutants of *Vibrio fischeri*. *J. Bacteriol.* 178:971–76
52. Latifi A, Foglino M, Tanaka K, Williams P, Lazdunski A. 1996. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhIR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol. Microbiol.* 21:1137–46
53. Lazazzera BA, Solomon JM, Grossman AD. 1997. An exported peptide functions intracellularly to contribute to cell density signaling in *B. subtilis*. *Cell* 89:917–25
54. Lee DH, Jeong HS, Jeong HG, Kim KM, Kim H, Choi SH. 2008. A consensus sequence for binding of SmcR, a *Vibrio vulnificus* LuxR homologue, and genome-wide identification of the SmcR regulon. *J. Biol. Chem.* 283:23610–18
55. Lee MS, Morrison DA. 1999. Identification of a new regulator in *Streptococcus pneumoniae* linking quorum sensing to competence for genetic transformation. *J. Bacteriol.* 181:5004–16
56. Lenz DH, Bassler BL. 2007. The small nucleoid protein Fis is involved in *Vibrio cholerae* quorum sensing. *Mol. Microbiol.* 63:859–71

57. Lenz DH, Mok KC, Lilley BN, Kulkarni RV, Wingreen NS, Bassler BL. 2004. The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in *Vibrio harveyi* and *Vibrio cholerae*. *Cell* 118:69–82
58. Lilley BN, Bassler BL. 2000. Regulation of quorum sensing in *Vibrio harveyi* by LuxO and sigma-54. *Mol. Microbiol.* 36:940–54
59. Lin W, Kovacicova G, Skorupski K. 2005. Requirements for *Vibrio cholerae* HapR binding and transcriptional repression at the *hapR* promoter are distinct from those at the *aphA* promoter. *J. Bacteriol.* 187:3013–19
60. Long T, Tu KC, Wang Y, Mehta P, Ong NP, et al. 2009. Quantifying the integration of quorum-sensing signals with single-cell resolution. *PLoS Biol.* 7:e68
61. Luo P, Li H, Morrison DA. 2003. ComX is a unique link between multiple quorum sensing outputs and competence in *Streptococcus pneumoniae*. *Mol. Microbiol.* 50:623–33
62. Luo P, Morrison DA. 2003. Transient association of an alternative sigma factor, ComX, with RNA polymerase during the period of competence for genetic transformation in *Streptococcus pneumoniae*. *J. Bacteriol.* 185:349–58
63. Luo ZQ, Su S, Farrand SK. 2003. In situ activation of the quorum-sensing transcription factor TraR by cognate and noncognate acyl-homoserine lactone ligands: kinetics and consequences. *J. Bacteriol.* 185:5665–72
64. Lupp C, Ruby EG. 2004. *Vibrio fischeri* LuxS and AinS: comparative study of two signal synthases. *J. Bacteriol.* 186:3873–81
65. Lupp C, Urbanowski M, Greenberg EP, Ruby EG. 2003. The *Vibrio fischeri* quorum-sensing systems *ain* and *lux* sequentially induce luminescence gene expression and are important for persistence in the squid host. *Mol. Microbiol.* 50:319–31
66. Lyon GJ, Wright JS, Christopoulos A, Novick RP, Muir TW. 2002. Reversible and specific extracellular antagonism of receptor-histidine kinase signaling. *J. Biol. Chem.* 277:6247–53
67. Lyon GJ, Wright JS, Muir TW, Novick RP. 2002. Key determinants of receptor activation in the *agr* autoinducing peptides of *Staphylococcus aureus*. *Biochemistry* 41:10095–104
68. Maddock JR, Shapiro L. 1993. Polar location of the chemoreceptor complex in the *Escherichia coli* cell. *Science* 259:1717–23
69. Magnuson R, Solomon J, Grossman AD. 1994. Biochemical and genetic characterization of a competence pheromone from *B. subtilis*. *Cell* 77:207–16
70. Mayville P, Ji G, Beavis R, Yang H, Goger M, et al. 1999. Structure-activity analysis of synthetic autoinducing thiolactone peptides from *Staphylococcus aureus* responsible for virulence. *Proc. Natl. Acad. Sci. USA* 96:1218–23
71. McGrath S, Wade DS, Pesci EC. 2004. Dueling quorum sensing systems in *Pseudomonas aeruginosa* control the production of the Pseudomonas quinolone signal (PQS). *FEMS Microbiol. Lett.* 230:27–34
72. Medina G, Juarez K, Diaz R, Soberon-Chavez G. 2003. Transcriptional regulation of *Pseudomonas aeruginosa* *rbIR*, encoding a quorum-sensing regulatory protein. *Microbiology* 149:3073–81
73. Meibom KL, Blokesch M, Dolganov NA, Wu CY, Schoolnik GK. 2005. Chitin induces natural competence in *Vibrio cholerae*. *Science* 310:1824–7
74. Meighen EA. 1991. Molecular biology of bacterial bioluminescence. *Microbiol. Rev.* 55:123–42
75. Miller MB, Skorupski K, Lenz DH, Taylor RK, Bassler BL. 2002. Parallel quorum sensing systems converge to regulate virulence in *Vibrio cholerae*. *Cell* 110:303–14
76. Miller ST, Xavier KB, Campagna SR, Taga ME, Semmelhack MF, et al. 2004. *Salmonella typhimurium* recognizes a chemically distinct form of the bacterial quorum-sensing signal AI-2. *Mol. Cell* 15:677–87
77. Milton DL. 2006. Quorum sensing in vibrios: complexity for diversification. *Int. J. Med. Microbiol.* 296:61–71
78. Milton DL, Chalker VJ, Kirke D, Hardman A, Camara M, Williams P. 2001. The LuxM homologue VanM from *Vibrio anguillarum* directs the synthesis of N-(3-hydroxyhexanoyl)homoserine lactone and N-hexanoylhomoserine lactone. *J. Bacteriol.* 183:3537–47
79. Minogue TD, Wehland-von Trebra M, Bernhardt F, von Bodman SB. 2002. The autoregulatory role of EsaR, a quorum-sensing regulator in *Pantoea stewartii* ssp. *stewartii*: evidence for a repressor function. *Mol. Microbiol.* 44:1625–35

80. Mok KC, Wingreen NS, Bassler BL. 2003. *Vibrio harveyi* quorum sensing: a coincidence detector for two autoinducers controls gene expression. *EMBO J.* 22:870–81
81. More MI, Finger LD, Stryker JL, Fuqua C, Eberhard A, Winans SC. 1996. Enzymatic synthesis of a quorum-sensing autoinducer through use of defined substrates. *Science* 272:1655–58
82. Nadell CD, Xavier JB, Foster KR. 2009. The sociobiology of biofilms. *FEMS Microbiol. Rev.* 33:206–24
83. Nadell CD, Xavier JB, Levin SA, Foster KR. 2008. The evolution of quorum sensing in bacterial biofilms. *PLoS Biol.* 6:e14
84. Neiditch MB, Federle MJ, Miller ST, Bassler BL, Hughson FM. 2005. Regulation of LuxPQ receptor activity by the quorum-sensing signal autoinducer-2. *Mol. Cell* 18:507–18
85. Neiditch MB, Federle MJ, Pompeani AJ, Kelly RC, Swem DL, et al. 2006. Ligand-induced asymmetry in histidine sensor kinase complex regulates quorum sensing. *Cell* 126:1095–108
86. Ninfa EG, Atkinson MR, Kamberov ES, Ninfa AJ. 1993. Mechanism of autophosphorylation of *Escherichia coli* nitrogen regulator II (NRII or NtrB): trans-phosphorylation between subunits. *J. Bacteriol.* 175:7024–32
87. Novick RP, Geisinger E. 2008. Quorum sensing in staphylococci. *Annu. Rev. Genet.* 42:541–64
88. Parsek MR, Greenberg EP. 2005. Sociomicrobiology: the connections between quorum sensing and biofilms. *Trends Microbiol.* 13:27–33
89. Passador L, Cook JM, Gambello MJ, Rust L, Iglewski BH. 1993. Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science* 260:1127–30
90. Pearson JP, Gray KM, Passador L, Tucker KD, Eberhard A, et al. 1994. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc. Natl. Acad. Sci. USA* 91:197–201
91. Pearson JP, Passador L, Iglewski BH, Greenberg EP. 1995. A second *N*-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* 92:1490–94
92. Pesci EC, Milbank JB, Pearson JP, McKnight S, Kende AS, et al. 1999. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* 96:11229–34
93. Pesci EC, Pearson JP, Seed PC, Iglewski BH. 1997. Regulation of las and rhl quorum sensing in *Pseudomonas aeruginosa*. *J. Bacteriol.* 179:3127–32
94. Pestova EV, Havarstein LS, Morrison DA. 1996. Regulation of competence for genetic transformation in *Streptococcus pneumoniae* by an auto-induced peptide pheromone and a two-component regulatory system. *Mol. Microbiol.* 21:853–62
95. Peterson S, Cline RT, Tettelin H, Sharov V, Morrison DA. 2000. Gene expression analysis of the *Streptococcus pneumoniae* competence regulons by use of DNA microarrays. *J. Bacteriol.* 182:6192–202
96. Piper KR, Beck von Bodman S, Farrand SK. 1993. Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction. *Nature* 362:448–50
97. Pompeani AJ, Irgon JJ, Berger MF, Bulyk ML, Wingreen NS, Bassler BL. 2008. The *Vibrio harveyi* master quorum-sensing regulator, LuxR, a TetR-type protein is both an activator and a repressor: DNA recognition and binding specificity at target promoters. *Mol. Microbiol.* 70:76–88
98. Rickard AH, Palmer RJ Jr, Blehert DS, Campagna SR, Semmelhack MF, et al. 2006. Autoinducer 2: a concentration-dependent signal for mutualistic bacterial biofilm growth. *Mol. Microbiol.* 60:1446–56
99. Rosenfeld N, Elowitz MB, Alon U. 2002. Negative autoregulation speeds the response times of transcription networks. *J. Mol. Biol.* 323:785–93
100. Ruby EG. 1996. Lessons from a cooperative, bacterial-animal association: the *Vibrio fischeri*-*Euprymna scolopes* light organ symbiosis. *Annu. Rev. Microbiol.* 50:591–624
101. Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS, West SA. 2009. Quorum sensing and the social evolution of bacterial virulence. *Curr. Biol.* 19:341–45
102. Sandoz KM, Mitzimberg SM, Schuster M. 2007. Social cheating in *Pseudomonas aeruginosa* quorum sensing. *Proc. Natl. Acad. Sci. USA* 104:15876–81
103. Schaefer AL, Greenberg EP, Oliver CM, Oda Y, Huang JJ, et al. 2008. A new class of homoserine lactone quorum-sensing signals. *Nature* 454:595–99
104. Schaefer AL, Val DL, Hanzelka BL, Cronan JE Jr, Greenberg EP. 1996. Generation of cell-to-cell signals in quorum sensing: acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. *Proc. Natl. Acad. Sci. USA* 93:9505–9

---

**103. First study to show that bacteria use the extracellular fatty acid pool (e.g., *p*-coumaric acid) for AHL biosynthesis.**

---



105. Schauder S, Shokat K, Surette MG, Bassler BL. 2001. The LuxS family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Mol. Microbiol.* 41:463–76
106. Schuster M, Greenberg EP. 2008. LuxR-type proteins in *Pseudomonas aeruginosa* quorum sensing: distinct mechanisms with global implications. In *Chemical Communication Among Bacteria*, ed. SC Winans, BL Bassler, pp. 133–44. Washington, DC: ASM Press
107. Schuster M, Urbanowski ML, Greenberg EP. 2004. Promoter specificity in *Pseudomonas aeruginosa* quorum sensing revealed by DNA binding of purified LasR. *Proc. Natl. Acad. Sci. USA* 101:15833–39
108. Seed PC, Passador L, Iglewski BH. 1995. Activation of the *Pseudomonas aeruginosa* lasI gene by LasR and the Pseudomonas autoinducer PAI: an autoinduction regulatory hierarchy. *J. Bacteriol.* 177:654–59
109. Simon MI, Crane BR, Crane A. 2007. *Two-component Signaling Systems*. San Diego: Academic
110. Solomon JM, Lazazzera BA, Grossman AD. 1996. Purification and characterization of an extracellular peptide factor that affects two different developmental pathways in *Bacillus subtilis*. *Genes Dev.* 10:2014–24
111. Solomon JM, Magnuson R, Srivastava A, Grossman AD. 1995. Convergent sensing pathways mediate response to two extracellular competence factors in *Bacillus subtilis*. *Genes Dev.* 9:547–58
112. Sourjik V, Berg HC. 2004. Functional interactions between receptors in bacterial chemotaxis. *Nature* 428:437–41
113. Spirig T, Tiaden A, Kiefer P, Buchrieser C, Vorholt JA, Hilbi H. 2008. The Legionella autoinducer synthase LqsA produces an alpha-hydroxyketone signaling molecule. *J. Biol. Chem.* 283:18113–23
114. Stevens AM, Dolan KM, Greenberg EP. 1994. Synergistic binding of the *Vibrio fischeri* LuxR transcriptional activator domain and RNA polymerase to the lux promoter region. *Proc. Natl. Acad. Sci. USA* 91:12619–23
115. Studdert CA, Parkinson JS. 2005. Insights into the organization and dynamics of bacterial chemoreceptor clusters through in vivo crosslinking studies. *Proc. Natl. Acad. Sci. USA* 102:15623–28
116. Sung CK, Morrison DA. 2005. Two distinct functions of ComW in stabilization and activation of the alternative sigma factor ComX in *Streptococcus pneumoniae*. *J. Bacteriol.* 187:3052–61
117. Surette MG, Miller MB, Bassler BL. 1999. Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: a new family of genes responsible for autoinducer production. *Proc. Natl. Acad. Sci. USA* 96:1639–44
118. Svenningsen SL, Tu KC, Bassler BL. 2009. Gene dosage compensation calibrates four regulatory RNAs to control *Vibrio cholerae* quorum sensing. *EMBO J.* 28:429–39
119. Svenningsen SL, Waters CM, Bassler BL. 2008. A negative feedback loop involving small RNAs accelerates *Vibrio cholerae*'s transition out of quorum-sensing mode. *Genes Dev.* 22:226–38
120. Swem LR, Swem DL, Wingreen NS, Bassler BL. 2008. Deducing receptor signaling parameters from in vivo analysis: LuxN/AI-1 quorum sensing in *Vibrio harveyi*. *Cell* 134:461–73
121. Sztajer H, Lemme A, Vilchez R, Schulz S, Geffers R, et al. 2008. Autoinducer-2-regulated genes in Streptococcus mutants UA159 and global metabolic effect of the luxS mutation. *J. Bacteriol.* 190:401–15
122. Tiaden A, Spirig T, Carranza P, Bruggemann H, Riedel K, et al. 2008. Synergistic contribution of the *Legionella pneumophila* lqs genes to pathogen-host interactions. *J. Bacteriol.* 190:7532–47
123. Tiaden A, Spirig T, Weber SS, Bruggemann H, Bosshard R, et al. 2007. The *Legionella pneumophila* response regulator LqsR promotes host cell interactions as an element of the virulence regulatory network controlled by RpoS and LetA. *Cell Microbiol.* 9:2903–20
124. Tu KC, Bassler BL. 2007. Multiple small RNAs act additively to integrate sensory information and control quorum sensing in *Vibrio harveyi*. *Genes Dev.* 21:221–33
- 124a. Tu KC, Long T, Svenningsen SL, Wingreen NS, Bassler BL. 2009. Synergistic negative feedback loops involving small regulatory RNAs precisely control the *Vibrio harveyi* quorum-sensing response. Submitted
125. Tu KC, Waters CM, Svenningsen SL, Bassler BL. 2008. A small-RNA-mediated negative feedback loop controls quorum-sensing dynamics in *Vibrio harveyi*. *Mol. Microbiol.* 70:896–907
126. Urbanowski ML, Lostroh CP, Greenberg EP. 2004. Reversible acyl-homoserine lactone binding to purified *Vibrio fischeri* LuxR protein. *J. Bacteriol.* 186:631–37
127. Vannini A, Volpari C, Gargioli C, Muraglia E, Cortese R, et al. 2002. The crystal structure of the quorum sensing protein TraR bound to its autoinducer and target DNA. *EMBO J.* 21:4393–401

---

120. Using mutagenesis, mathematical modeling, and antagonist analyses, signaling parameters of the membrane-bound quorum-sensing receptor LuxN in *V. harveyi* were determined

---

128. Vendeville A, Winzer K, Heurlier K, Tang CM, Hardie KR. 2005. Making 'sense' of metabolism: autoinducer-2, LuxS and pathogenic bacteria. *Nat. Rev. Microbiol.* 3:383–96
129. von Bodman SB, Ball JK, Faini MA, Herrera CM, Minogue TD, et al. 2003. The quorum sensing negative regulators EsaR and ExpR(Ecc), homologues within the LuxR family, retain the ability to function as activators of transcription. *J. Bacteriol.* 185:7001–7
130. von Bodman SB, Farrand SK. 1995. Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia stewartii* require induction by an *N*-acylhomoserine lactone autoinducer. *J. Bacteriol.* 177:5000–8
131. von Bodman SB, Majerczak DR, Coplin DL. 1998. A negative regulator mediates quorum-sensing control of exopolysaccharide production in *Pantoea stewartii* subsp. *stewartii*. *Proc. Natl. Acad. Sci. USA* 95:7687–92
132. Wade DS, Calfee MW, Rocha ER, Ling EA, Engstrom E, et al. 2005. Regulation of *Pseudomonas* quinolone signal synthesis in *Pseudomonas aeruginosa*. *J. Bacteriol.* 187:4372–80
133. Waters CM, Bassler BL. 2005. Quorum sensing: cell-to-cell communication in bacteria. *Annu. Rev. Cell Dev. Biol.* 21:319–46
134. Waters CM, Bassler BL. 2006. The *Vibrio harveyi* quorum-sensing system uses shared regulatory components to discriminate between multiple autoinducers. *Genes Dev.* 20:2754–67
135. Waters LS, Storz G. 2009. Regulatory RNAs in bacteria. *Cell* 136:615–28
136. Watson WT, Minogue TD, Val DL, von Bodman SB, Churchill ME. 2002. Structural basis and specificity of acyl-homoserine lactone signal production in bacterial quorum sensing. *Mol. Cell* 9:685–94
137. West SA, Griffin AS, Gardner A, Diggle SP. 2006. Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* 4:597–607
138. Williams P, Winzer K, Chan WC, Camara M. 2007. Look who's talking: communication and quorum sensing in the bacterial world. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 362:1119–34
139. Winson MK, Camara M, Latifi A, Foglino M, Chhabra SR, et al. 1995. Multiple *N*-acyl-L-homoserine lactone signal molecules regulate production of virulence determinants and secondary metabolites in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* 92:9427–31
140. Wolfgang MC, Lee VT, Gilmore ME, Lory S. 2003. Coordinate regulation of bacterial virulence genes by a novel adenylate cyclase-dependent signaling pathway. *Dev. Cell* 4:253–63
141. Wright JS 3rd, Lyon GJ, George EA, Muir TW, Novick RP. 2004. Hydrophobic interactions drive ligand-receptor recognition for activation and inhibition of staphylococcal quorum sensing. *Proc. Natl. Acad. Sci. USA* 101:16168–73
142. Xavier KB, Bassler BL. 2003. LuxS quorum sensing: more than just a numbers game. *Curr. Opin. Microbiol.* 6:191–97
143. Yang Y, Inouye M. 1991. Intermolecular complementation between two defective mutant signal-transducing receptors of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 88:11057–61
144. Yao Y, Martinez-Yamout MA, Dickerson TJ, Brogan AP, Wright PE, Dyson HJ. 2006. Structure of the *Escherichia coli* quorum sensing protein SdiA: activation of the folding switch by acyl homoserine lactones. *J. Mol. Biol.* 355:262–73
145. Zhang RG, Pappas T, Brace JL, Miller PC, Oulmassov T, et al. 2002. Structure of a bacterial quorum-sensing transcription factor complexed with pheromone and DNA. *Nature* 417:971–74
146. Zhu J, Mekalanos JJ. 2003. Quorum sensing-dependent biofilms enhance colonization in *Vibrio cholerae*. *Dev. Cell* 5:647–56
147. Zhu J, Miller MB, Vance RE, Dziejman M, Bassler BL, Mekalanos JJ. 2002. Quorum-sensing regulators control virulence gene expression in *Vibrio cholerae*. *Proc. Natl. Acad. Sci. USA* 99:3129–34
148. Zhu J, Winans SC. 1999. Autoinducer binding by the quorum-sensing regulator TraR increases affinity for target promoters in vitro and decreases TraR turnover rates in whole cells. *Proc. Natl. Acad. Sci. USA* 96:4832–37
149. Zhu J, Winans SC. 2001. The quorum-sensing transcriptional regulator TraR requires its cognate signaling ligand for protein folding, protease resistance, and dimerization. *Proc. Natl. Acad. Sci. USA* 98:1507–12



# Contents

Genetic and Epigenetic Mechanisms Underlying Cell-Surface Variability in Protozoa and Fungi <i>Kevin J. Verstrepen and Gerald R. Fink</i> .....	1
Regressive Evolution in <i>Astyanax</i> Cavefish <i>William R. Jeffery</i> .....	25
Mimivirus and its Virophage <i>Jean-Michel Claverie and Chantal Abergel</i> .....	49
Regulation Mechanisms and Signaling Pathways of Autophagy <i>Congcong He and Daniel J. Klionsky</i> .....	67
The Role of Mitochondria in Apoptosis <i>Chunxin Wang and Richard J. Youle</i> .....	95
Biom mineralization in Humans: Making the Hard Choices in Life <i>Kenneth M. Weiss, Kazubiko Kawasaki, and Anne V. Buchanan</i> .....	119
Active DNA Demethylation Mediated by DNA Glycosylases <i>Jian-Kang Zhu</i> .....	143
Gene Amplification and Adaptive Evolution in Bacteria <i>Dan I. Andersson and Diarmaid Hughes</i> .....	167
Bacterial Quorum-Sensing Network Architectures <i>Wai-Leung Ng and Bonnie L. Bassler</i> .....	197
How the Fanconi Anemia Pathway Guards the Genome <i>George-Lucian Moldovan and Alan D. D'Andrea</i> .....	223
Nucleomorph Genomes <i>Christa Moore and John M. Archibald</i> .....	251
Mechanism of Auxin-Regulated Gene Expression in Plants <i>Elisabeth J. Chapman and Mark Estelle</i> .....	265
Maize Centromeres: Structure, Function, Epigenetics <i>James A. Birchler and Fangpu Han</i> .....	287

The Functional Annotation of Mammalian Genomes: The Challenge of Phenotyping <i>Steve D.M. Brown, Wolfgang Wurst, Ralf Kühn, and John Hancock</i> .....	305
Thioredoxins and Glutaredoxins: Unifying Elements in Redox Biology <i>Yves Meyer, Bob B. Buchanan, Florence Vignols, and Jean-Philippe Reichheld</i> .....	335
Roles for BMP4 and CAM1 in Shaping the Jaw: Evo-Devo and Beyond <i>Kevin J. Parsons and R. Craig Albertson</i> .....	369
Regulation of Tissue Growth through Nutrient Sensing <i>Ville Hietakangas and Stephen M. Cohen</i> .....	389
Hearing Loss: Mechanisms Revealed by Genetics and Cell Biology <i>Aniel A. Dror and Karen B. Avraham</i> .....	411
The Kinetochore and the Centromere: A Working Long Distance Relationship <i>Marcin R. Przewłoka and David M. Glover</i> .....	439
Multiple Roles for Heterochromatin Protein 1 Genes in <i>Drosophila</i> <i>Danielle Vermaak and Harmit S. Malik</i> .....	467
Genetic Control of Programmed Cell Death During Animal Development <i>Barbara Conradt</i> .....	493
Cohesin: Its Roles and Mechanisms <i>Kim Nasmyth and Christian H. Haering</i> .....	525
Histones: Annotating Chromatin <i>Eric I. Campos and Danny Reinberg</i> .....	559
Systematic Mapping of Genetic Interaction Networks <i>Scott J. Dixon, Michael Costanzo, Charles Boone, Brenda Andrews, and Anastasia Baryshnikova</i> .....	601

## Errata

An online log of corrections to *Annual Review of Genetics* articles may be found at <http://genet.annualreviews.org/errata.shtml>